

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 July 2003 (24.07.2003)

PCT

(10) International Publication Number
WO 03/059881 A2

(51) International Patent Classification⁷: C07D 207/34,
417/14, 417/12, 403/14, 409/14, A61K 31/40, 31/427,
31/404, 31/4025, A61P 31/00

Street, Thomas Graham Building, University of Strathclyde, Glasgow G1 1XL (GB).

(21) International Application Number: PCT/GB02/05916

(74) Agent: MCNEENEY, Stephen, P.; Eric Potter Clarkson, Park View House, 58 The Ropewalk, Nottingham NG1 5DD (GB).

(22) International Filing Date:
24 December 2002 (24.12.2002)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0130868.3 24 December 2001 (24.12.2001) GB

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): UNIVERSITY OF STRATHCLYDE [GB/GB]; 16 Richmond Street, Glasgow G1 1XQ (GB).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): KHALAF, Abedawn [GB/GB]; Department of Pure and Applied Chemistry, 295 Cathedral Street, Thomas Graham Building, University of Strathclyde, Glasgow G1 1XL (GB). WAIGH, Roger [GB/GB]; Department of Pharmaceutical Sciences, University of Strathclyde, 27 Taylor Street, Glasgow G4 0NR (GB). SUCKLING, Colin [GB/GB]; Department of Pure and Applied Chemistry, 295 Cathedral

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DNA MINOR GROOVE BINDING COMPOUNDS

(57) Abstract: There is provided an oligopeptide compound comprising: (a) at least one nitrogen-containing basic group attached to at least one end of the oligopeptide; and (b) two or more heterocyclic monomers, at least one of which is substituted in the heterocyclic part by a branched, cyclic or part cyclic C₃₋₅ alkyl group, or a pharmaceutically acceptable salt or solvate thereof; which compound, salt or solvate binds to the minor groove of DNA.



WO 03/059881 A2

DNA MINOR GROOVE BINDING COMPOUNDS

Field of the Invention

- 5 This invention relates to synthetic compounds that have affinity for nucleic acids, and in particular to compounds that bind to the minor groove of DNA.

Background and Prior Art

10

Because of its fundamental role in molecular biological processes, DNA represents an important target for drug action. Compounds that can recognise defined sequences of DNA have a wide variety of potential uses, such as the modulation of gene expression.

15

The outer surface of double-helical DNA has two channels, namely the major and minor grooves. Both of these grooves contain chemical information by way of arrangements of hydrogen-bond donors and acceptors, electrostatic charges, dipoles, hydrophobic regions and so on.

20

The major groove contains approximately twice the information content of the minor groove in terms of the number of potential hydrogen-bonding contacts. In view of this, the major groove is the preferred recognition site for cellular proteins such as control proteins, promoters and repressors.

25

In contrast, the minor groove is normally (with a few exceptions) relatively unoccupied. The vulnerability of the minor groove makes it a particularly useful target for compounds that bind to DNA. Indeed, perhaps for this very reason, the minor groove is the binding site for certain naturally occurring antibiotics (such as netropsin and distamycin).

30

Netropsin and distamycin are oligopeptides based on pyrrole amino acid monomers. These compounds both bind to DNA with dissociation constants in the order of 10^{-5} M. They also show a preference for AT-rich regions of DNA. Although they have intrinsic biological activity, netropsin and distamycin also have many limitations including toxicity, moderate affinity and limited selectivity. A number of workers have therefore prepared synthetic analogues of netropsin and distamycin, with a view to overcoming these disadvantages. Many of these compounds are reviewed by Sondhi *et al.* (*Curr. Med. Chem.* 4, 313 (1997)), Reddy *et al.* (10 *Pharmacology & Therapeutics* 84, 1 (1999)), Wemmer (*Biopolymers* 52, 197 (2001)) and Dervan (*Bioorg. Med. Chem.* 9, 2215 (2001)).

Compounds designed to bind to DNA regions containing GC base pairs are described in, for example: *Anti-Cancer Drug Design* 5, 3 (1990); *Proc. Natl. Acad. Sci. USA* 89, 7586 (1992); *Biochemistry* 32, 4237 (1993); 15 *Science* 266, 647 (1994); *Anti-Cancer Drug Design* 10, 155 (1995); *Bioorg. Med. Chem.* 8, 985 (2000); and *Mol. Biol.* 34, 357 (2000). Various other netropsin and distamycin analogues are described in: *J. Am. Chem. Soc.* 114(15), 5911 (1992); *Biochemistry* 31, 8349 (1992); *Bioconjugate Chem.* 20 5, 475 (1994); *Biochem. Biophys. Res. Commun.* 222, 764 (1996); *J. Med. Chem.* 43, 3257 (2000); and *Tetrahedron* 56, 5225 (2000). Further, the use of certain netropsin and distamycin analogues as antimicrobial, antiviral and/or antitumor agents is described in *Molecular Pharmacology* 54, 280 (1998), *Bioorg. Med. Chem. Lett.* 6(18), 2169 (1996), *J. Med. Chem.* 45, 25 805 (2002), *Bioorg. Med. Chem. Lett.* 12, 2007 (2002), international patent applications WO 97/28123, WO 98/21202, WO 01/74898 and WO 02/00650, as well as in US patent numbers 4,912,199, 5,273,991, 5,637,621, 5,698,674 and 5,753,629. Methods of synthesising analogues of netropsin and distamycin are described in US 6,090,947.

Cellular uptake of distamycin analogues is described in *Bioorg. Med. Chem. Lett.* 11, 769 (2001).

None of the above-mentioned documents describe oligopeptide analogues of netropsin or distamycin, which analogues comprise at least two heterocyclic monomers, at least one of which is substituted in the heterocyclic part by a branched, cyclic or part cyclic C₃₋₅ alkyl group. Surprisingly, we have found that compounds of this type bind with a high affinity and specificity to the minor groove of DNA.

10

Description of the Invention

According to the invention there is provided oligopeptide compounds comprising:

- 15 (a) at least one nitrogen-containing basic group attached to at least one end of the oligopeptide; and
- (b) two or more heterocyclic monomers, at least one of which is substituted in the heterocyclic part by a branched, cyclic or part cyclic C₃₋₅ alkyl group,

20 or a pharmaceutically acceptable salt or solvate thereof;

which compounds, salts or solvates bind to the minor groove of DNA, and which compounds are referred to hereinafter as "the compounds of the invention".

25 When used herein, the term "oligopeptide" includes organometallic and, particularly, organic compounds that contain at least one amide (e.g. -C(O)NH-) group that links different parts of the molecule. Preferred oligopeptide compounds are those of low molecular weight (e.g. below 2000, such as below 1500 and, particularly, below 1000 g mol⁻¹). Preferred

30 oligopeptide compounds also include those that comprise heterocyclic

amino acids (i.e. heterocyclic groups that possess both carboxylic acid and amine functionalities) that are coupled to each other *via* linkages that include at least one amide bond (e.g. a direct amide bond or a linkage formed by reaction of two heterocyclic amino acids with a compound that possesses two carboxylic acid groups, two amine groups or one acid and one amine group). Suitable compounds possessing two carboxylic acid groups, two amine groups or one acid and one amine group are known to those skilled in the art, for example the "linkers" disclosed in *Curr. Med. Chem.* 4, 313-358 (1997), *Pharmacology & Therapeutics* 84, 1-111 (1999), *J. Med. Chem.* 43, 3257-3266 (2000), *Tetrahedron* 56, 5225-5239 (2000), the disclosures of which documents are hereby incorporated by reference.

When used herein, the term "nitrogen-containing basic group" includes nitrogen-containing compounds that form salts with organic or inorganic acids. As such, the term includes nitrogen-containing groups that, in their neutral state, have a pK_a in water of at least 4 (preferably at least 5, 6 or 7, and particularly in the range of 8 to 12). The term therefore specifically includes amidino, guanidino and, particularly, amino groups, any of which may be cyclic or acyclic. Preferred nitrogen-containing basic groups include $-N(H)C(=NH)NH_2$, $-C(=NH)NH_2$ and, particularly, amino groups such as $-NH_2$, alkylamino (e.g. $-N(H)CH_3$) and dialkylamino (e.g. $-N(CH_3)_2$) groups and saturated azaheterocyclic groups such as pyrrolidinyl and piperazinyl. Particularly preferred nitrogen-containing basic groups include $-N(CH_3)_2$. For the avoidance of doubt, the nitrogen-containing basic group is a separate moiety from the two or more heterocyclic groups mentioned at point (b) above.

When used herein, the term "heterocyclic group" includes 4- to 12-membered (e.g. 5- to 10-membered) heterocyclic groups containing one or more heteroatoms selected from N, O and S. The term therefore includes

such groups that are mono- or bicyclic, and which may be saturated, part-unsaturated, aromatic or, where appropriate, part-aromatic. Preferred heterocyclic groups include aromatic or part-aromatic groups such as pyrrolyl, imidazolyl, thiazolyl, oxazolyl, benzoxazolyl, furanyl, thienyl, pyridyl and coumarinyl. Particularly preferred heterocyclic groups include pyrrolyl, imidazolyl, thiazolyl and oxazolyl.

By “substituted in the heterocyclic part”, we mean that each of the essential branched, cyclic or part cyclic C₃₋₅ alkyl groups is a direct substituent on the heterocyclic ring (whether attached to the ring *via* a heteroatom or otherwise) of each heterocyclic monomer bearing such a group.

In one embodiment of the invention, at least one heterocyclic group that is present (e.g. an imidazolyl, a thienyl or, particularly, a thiazolyl group) is substituted at a ring C-atom by the branched, cyclic or part cyclic C₃₋₅ alkyl group.

In another embodiment of the invention, at least one heterocyclic group that is present is a thiazolyl group (e.g. a 1,3-thiazolyl group that is substituted in the 5-position by the branched, cyclic or part cyclic C₃₋₅ alkyl group).

The compounds of the invention may be provided in a form rendering them bioavailable. When used herein, the term “bioavailable” includes compounds that, following administration, are in a form in which they can interact with a biological system, thereby providing a measurable therapeutic response. The term may thus be understood to include compounds that are provided to DNA in a form and/or level that is sufficient to provide a measurable desired or required therapeutic response.

Bioavailability of a compound may be predicted by a number of means known to those skilled in the art, including a measurement of a partition coefficient of the compound between water (for example at a pH of between 5 and 9) and an organic, water-immiscible solvent (e.g. octanol), which measurement can be used to predict the behaviour in body tissues of the compound in question (for a discussion of which see *J. Med. Chem.* 43, 3257-3266 (2000)).

Bioavailability may be achieved by providing compounds of the invention in a form (e.g. a pharmaceutical formulation) in which they are presented to DNA at an appropriate concentration to provide a measurable therapeutic response. Bioavailability may alternatively be achieved by changing the physicochemical properties of the active species, for example by improving water solubility by using techniques known to those skilled in the art (e.g. by the introduction of additional basic groups, such as described in *J. Med. Chem.* 43, 3257-3266 (2000)).

The compounds of the invention may have a high affinity for at least one DNA sequence. When used herein, the term "high affinity for at least one DNA sequence" includes compounds that, when bound to a minor groove of at least one DNA oligomer or polymer, have a dissociation constant of less than 10^{-5} M, preferably less than 10^{-6} M (such as 10^{-7} M) and particularly less than 10^{-8} M. In this respect, dissociation constants may be measured under conditions known to those skilled in the art, for example in water at room temperature (e.g. at or around 20°C) in the presence of a buffer (e.g. a buffer that stabilises the pH at 7.5, such as a borate (e.g. at 0.02M) or Tris/HCl (e.g. at 0.01 M) buffer) and at a DNA concentration of between 10 and 30 μ M (e.g. 20 μ M). Alternatively, dissociation constants may be estimated by a comparison of the binding affinity of a compound to a set

DNA sequence with the binding affinity of a well-known compound (e.g. distamycin) to that same sequence.

Unless otherwise specified, the term "DNA" refers to double-stranded DNA. Further, when used herein, the term "DNA sequence" includes any part of (or the whole of) a DNA oligomer or polymer spanning three or more base pairs.


Preferred compounds of the invention also include those in which each essential branched, cyclic or part cyclic C₃₋₅ alkyl substituent on a heterocyclic monomer is:

other than cyclopropyl;

a branched C₃₋₅ alkyl group such as cyclopropylmethyl, cyclopentyl, isopentyl or isopropyl (e.g. isopentyl and, particularly, isopropyl).

15

According to a further aspect of the invention, there is provided compounds of formula I,



wherein

20 R^1 represents Het¹, R^{1a}C(O)- or D-A-N(H)-[Q]_n-C(O)-E-C(O)-;

R^{1a} represents

H,

aryl (which latter group is optionally substituted by one or more substituents selected from OH, halo, cyano, nitro, N(R^{3a})R^{3b}, C₁₋₄ alkyl and C₁₋₄ alkoxy),

25

aromatic or part-aromatic C₁₃₋₁₄ tricyclic carbocyclyl (which latter group is optionally substituted by one or more substituents selected from OH, halo, cyano, nitro, N(R^{3a})R^{3b}, C₁₋₄ alkyl and C₁₋₄ alkoxy, and

which latter group, if part-aromatic, is optionally substituted in the non-aromatic part by one or two oxo groups) or

C₁₋₁₂ alkyl (which latter group is optionally substituted and/or terminated by one or more substituents selected from halo and aryl (which latter group is optionally substituted by one or more substituents selected from OH, halo, cyano, nitro, N(R^{3a})R^{3b}, C₁₋₄ alkyl and C₁₋₄ alkoxy));

A represents, at each occurrence when used herein, C₂₋₆ alkylene or A¹-C(O)N(H)-A², wherein A² is attached to the group D;

A¹ represents C₁₋₄ alkylene;

A² represents C₂₋₅ alkylene;

D represents, at each occurrence when used herein, -N(R^{2a})R^{2b}, -C(=NR^{2c})N(R^{2d})R^{2e} or -N(R^{2f})C(=NR^{2g})N(H)R^{2h};


R^{2a} and R^{2b} independently represent H, C₁₋₆ alkyl, Het² or R^{2a} and R^{2b} together represent (CH₂)₃₋₆, which alkylene group is optionally interrupted by NR⁴ and/or is optionally substituted by one or more C₁₋₄ alkyl groups;

R⁴ represents H, C₁₋₆ alkyl or Het³;

R^{2c} to R^{2h} independently represent H or C₁₋₆ alkyl;

20

E represents -E¹-Het⁴-, E^{2a}-, -(CH₂)₀₋₃N(H)C(O)-E^{2b}-C(O)N(H)(CH₂)₀₋₃- or a structural fragment of the formula



wherein E³ represents (CH₂)₁₋₂, CH=CH, CH=N, CH₂-N(R^a), (CH₂)₀₋₁C(O), (CH₂)₀₋₁O or (CH₂)₀₋₁S;

R^a represents H or C₁₋₆ alkyl;

E¹ represents (CH₂)₀₋₂ or CH=CH;

25


E^{2a} and E^{2b} independently represent C_{2-4} alkenylene, C_{3-6} cycloalkylene, phenylene or naphthylene;

Het¹ to Het⁴ independently represent four- to twelve-membered heterocyclic groups containing one or more heteroatoms selected from N, O and S, which heterocyclic groups are optionally substituted by one or more substituents selected from =O, OH, halo, cyano, nitro, $N(R^{3a})R^{3b}$, C_{1-4} alkyl and C_{1-4} alkoxy;


R^{3a} and R^{3b} independently represent, at each occurrence when used herein, H or C_{1-4} alkyl, or R^{3a} represents $-C(O)R^5$; R^5 represents H or C_{1-4} alkyl;

n represents, at each occurrence when used herein, 2, 3, 4 or 5;


each individual Q independently represents a structural fragment of formula Ia, Ib, Ic, Id, Ie or If



10



1e



1f

wherein

R^6 represents H or C_{1-6} alkyl;

R^7 represents C_{1-12} alkyl;

5 R^8 , R^9 , R^{10} and R^{11} independently represent H or C_{1-12} alkyl;

G represents CH or N;

L represents O or S;

p, q and r independently represent 0, 1, 2 or 3; and

10 provided that the compound comprises at least one structural fragment of formula Ib, Ic, Id, 1e or 1f in which R^6 or R^7 , R^8 , R^9 , R^{10} or R^{11} , respectively, represents branched, cyclic or part cyclic C_{3-5} alkyl;

or a pharmaceutically acceptable derivative thereof,

15

which compounds are also referred to hereinafter as “the compounds of the invention”.

Unless otherwise specified, alkyl groups and alkoxy groups as defined
 20 herein may be straight-chain or, when there is a sufficient number (i.e. a minimum of three) of carbon atoms, be branched-chain and/or cyclic. Further, when there is a sufficient number (i.e. a minimum of four) of carbon atoms, such alkyl and alkoxy groups may also be part cyclic/acyclic. Such alkyl and alkoxy groups may also be saturated or, when there is a

sufficient number (i.e. a minimum of two) of carbon atoms, be unsaturated and/or interrupted by one or more oxygen and/or sulfur atoms. Unless otherwise specified, alkyl and alkoxy groups may also be substituted by one or more halo, and especially fluoro, atoms.

5

Unless otherwise specified, alkylene groups as defined herein may be straight-chain or, when there is a sufficient number (i.e. a minimum of two) of carbon atoms, be branched-chain. Such alkylene chains may also be saturated or, when there is a sufficient number (i.e. a minimum of two) of
10 carbon atoms, be unsaturated and/or interrupted by one or more oxygen and/or sulfur atoms. Unless otherwise specified, alkylene groups may also be substituted by one or more halo atoms.

The term "aryl", when used herein, includes C₆₋₁₀ aryl groups such as phenyl, naphthyl and the like. When substituted, aryl groups are preferably
15 substituted by between one and three substituents.

The term "aromatic or part-aromatic C₁₃₋₁₄ tricyclic carbocyclyl", when used herein includes fluorenyl, anthracenyl, 9,10-dihydroanthracenyl, phenanthrenyl, 9,10-dihydrophenanthrenyl and the like.
20

The term "halo", when used herein, includes fluoro, chloro, bromo and iodo.

Het (Het¹ to Het⁴) groups that may be mentioned include those containing 1
25 to 4 heteroatoms (selected from the group oxygen, nitrogen and/or sulfur) and in which the total number of atoms in the ring system are between five and twelve. Het (Het¹ to Het⁴) groups may be fully saturated, partly unsaturated, wholly aromatic, partly aromatic and/or bicyclic in character. Heterocyclic groups that may be mentioned include benzodioxanyl,
30 benzodioxepanyl, benzodioxolyl, benzofuranyl, benzofurazanyl,

benzimidazolyl, benzomorpholinyl, benzothiazolyl, benzothiophenyl, benzoxazolyl, chromanyl, cinnolinyl, coumarinyl, dioxanyl, furanyl, hydantoinyl, imidazolyl, imidazo[1,2-*a*]pyridinyl, indolyl, isoquinolinyl, isoxazolyl, maleimido, morpholinyl, oxazolyl, phthalazinyl, piperazinyl, 5 piperidinyl, purinyl, pyranyl, pyrazinyl, pyrazolyl, pyridinyl, pyrimidinyl, pyrrolidinonyl, pyrrolidinyl, pyrrolinyl, pyrrolyl, quinazolinyl, quinolinyl, 3-sulfolenyl, tetrahydropyranyl, tetrahydrofuranyl, thiazolyl, thienyl, thiochromanyl, triazolyl and the like. Values of Het¹, Het² and Het³ that may be mentioned include benzoxazolyl. Values of Het⁴ that may be 10 mentioned include indolyl.

Pharmaceutically acceptable derivatives include salts and solvates. Salts which may be mentioned include acid addition salts.

15 Compounds of formula I may exhibit tautomerism. All tautomeric forms and mixtures thereof are included within the scope of the invention.

Compounds of formula I may also contain one or more asymmetric carbon atoms and may therefore exhibit optical and/or diastereoisomerism. 20 Diastereoisomers may be separated using conventional techniques, e.g. chromatography or fractional crystallisation. The various stereoisomers may be isolated by separation of a racemic or other mixture of the compounds using conventional, e.g. fractional crystallisation or HPLC, techniques. Alternatively the desired optical isomers may be made by 25 reaction of the appropriate optically active starting materials under conditions which will not cause racemisation or epimerisation, or by derivatisation, for example with a homochiral acid followed by separation of the diastereomeric esters by conventional means (e.g. HPLC, chromatography over silica). All stereoisomers are included within the 30 scope of the invention.

Compounds of formula I that may be mentioned include those in which:

R^{1a} represents H or C_{1-12} alkyl, which latter group is optionally substituted and/or terminated by one or more substituents selected from halo and aryl, which latter group is optionally substituted by one or more substituents selected from OH, halo, cyano, nitro, $N(R^{3a})R^{3b}$, C_{1-4} alkyl and C_{1-4} alkoxy;

the compound comprises at least one structural fragment of formula Ib, Ic, Id, Ie or If in which R^7 , R^8 , R^9 , R^{10} or R^{11} , respectively, represents branched, cyclic or part cyclic C_{3-5} alkyl.

10

Branched, cyclic or part cyclic C_{3-5} alkyl groups that may be mentioned include isopropyl, cyclopropylmethyl, isopentyl and cyclopentyl.

Preferred compounds of formula I include those in which:

15 R^{1a} represents

H,

phenyl, anthracenyl (which latter two groups are optionally substituted by one to three substituents selected from OH, halo, C_{1-4} alkyl and C_{1-4} alkoxy),

20 9,10-dihydroanthracenyl (which latter group is optionally substituted by one or two oxo groups and/or one to three substituents selected from OH, halo, C_{1-4} alkyl and C_{1-4} alkoxy) or

optionally unsaturated C_{1-8} alkyl, which latter group is substituted by one or more halo groups and/or terminated by phenyl, which latter

25 group is optionally substituted by one or more substituents selected from OH, halo, cyano, nitro, $N(R^{3a})R^{3b}$, C_{1-2} alkyl and C_{1-2} alkoxy;

A represents optionally unsaturated C_{2-5} alkylene or $A^1-C(O)N(H)-A^2$;

A^1 represents saturated C_{1-3} alkylene;


A^2 represents saturated C_{2-4} alkylene;

R^{2a} and R^{2b} independently represent H, C_{1-4} alkyl, Het² or R^{2a} and R^{2b} together represent $(CH_2)_{3-6}$, which alkylene group is optionally interrupted by NR^4 and/or is optionally substituted by a C_{1-2} alkyl group;

R^4 represents C_{1-4} alkyl or Het³;

5 R^{2c} to R^{2h} independently represent H;

E represents $-E^1-Het^4-$, *cis*- or *trans*-ethenylene, cyclopropylene, 1,3- or 1,4-phenylene, $-(CH_2)_{1-2}N(H)C(O)-phenylene-C(O)N(H)(CH_2)_{1-2}-$ or a structural fragment of the formula



10 wherein E^3 represents $(CH_2)_2$, $CH=CH$, $C(O)$ or S;

Het¹ to Het³ independently represent six- to ten-membered heterocyclic groups containing one or more heteroatoms selected from N, O and S, which heterocyclic groups are optionally substituted by one or more substituents selected from =O, OH, halo, cyano, nitro, $N(R^{3a})R^{3b}$, C_{1-2} alkyl

15 and C_{1-2} alkoxy;

Het⁴ represents an unsubstituted nine- or ten-membered aromatic heterocyclic group containing one or two N-atoms (e.g. indolyl, such as 2,5- or 2,6-indolyl);

R^{3a} and R^{3b} independently represent, at each occurrence when used herein,

20 H or C_{1-2} alkyl, or R^{3a} represents $-C(O)R^5$;

R^5 represents H or C_{1-2} alkyl;

R^6 represents H or C_{1-2} alkyl or, when G represents N, R^6 may also represent branched, cyclic or part cyclic C_{3-5} alkyl;

R^7 represents C_{1-8} alkyl;

25 R^8 , R^9 , R^{10} and R^{11} independently represent H or C_{1-8} alkyl;

p, q and r independently represent 0, 1 or 2;

n represents 3, 4 or 5 or, when R¹ represents methyl, Het¹ or D-A-N(H)-[Q]_n-C(O)-E-C(O)-, then n may also represent 2;

the compound comprises at least one structural fragment of formula Ib, Ic, Id, Ie or If in which R⁷, R⁸, R⁹, R¹⁰ or R¹¹, respectively, represents branched C₃₋₅ alkyl.

More preferred compounds of formula I include those in which:

R^{1a} represents

H

- 10 phenyl (which latter group is optionally substituted by C₁₋₂ alkoxy),
9,10-dioxo-9,10-dihydroanthracenyl (which latter group is optionally substituted by C₁₋₂ alkoxy),
saturated, optionally branched C₁₋₆ alkyl or
saturated C₁₋₃ *n*-alkyl, which latter group is terminated by phenyl
15 (which latter group is optionally substituted by C₁₋₂ alkoxy);

A represents saturated C₂₋₄ alkylene or A¹-C(O)N(H)-A²;

A¹ represents (CH₂)₁₋₃;

A² represents (CH₂)₂₋₄;


D represents -N(R^{2a})R^{2b};

- 20 R^{2a} and R^{2b} independently represent C₁₋₃ alkyl or Het², or R^{2a} and R^{2b} together represent (CH₂)₃₋₅, which alkylene group is optionally interrupted by NR⁴;

R⁴ represents C₁₋₃ alkyl or Het³;

- E represents -(2,5-indolyl)-, -(CH₂)₀₋₂-(2,6-indolyl)-, -CH=CH-(2,6-indolyl)-, *trans*-ethenylene, *trans*-cyclopropylene, 1,3- or 1,4-phenylene,
25 -CH₂N(H)C(O)-(1,3- or 1,4-phenylene)-C(O)N(H)CH₂- or one of the following structural fragments

16



- Het¹, Het² and/or Het³ independently represent nine- or ten-membered aromatic heterocycles containing one to three heteroatoms selected from N, O and S;
- R⁶ represents H or, when G represents N, R⁶ may also represent branched, cyclic or part cyclic C₃₋₅ alkyl;
- R⁷, R⁸, R⁹, R¹⁰ and R¹¹ independently represent saturated, optionally branched C₁₋₆ alkyl or R⁸ represents H;
- p, q and r independently represent 0 or 1;
- n represents 3 or 4, or, when R¹ represents methyl or D-A-N(H)-[Q]_n-C(O)-E-C(O)-, then n may also represent 2;
- the compound comprises at least one structural fragment of formula Ib, Ic, Id, Ie or If in which R⁷, R⁸, R⁹, R¹⁰ or R¹¹, respectively, represents iso- or sec-butyl, isopentyl or, particularly, isopropyl;
- the compound comprises at least one structural fragment of formula Ib, Ic, Id, Ie or If in which R⁷, R⁸, R⁹, R¹⁰ or R¹¹, respectively, represents ethyl or, particularly, methyl.

- Compounds of formula I that are more preferred still include those in which:

R^{1a} represents

H

- methoxyphenyl (e.g. *p*- or, particularly, *m*-methoxyphenyl),
9,10-dioxo-9,10-dihydroanthracenyl,
saturated C₁₋₃ alkyl (e.g. methyl) or
methoxyphenylacetyl (e.g. *m*- or, particularly, *p*-methoxyphenyl);
- 5 Het¹ represents a nine-membered aromatic heterocycle containing two heteroatoms selected from N, O and S (e.g. 1,3-benzoxazol-2-yl);
A represents saturated C₂₋₄ *n*-alkylene (e.g. *n*-propylene) or A¹-C(O)N(H)-A²;
A¹ represents (CH₂)₂;
- 10 A² represents (CH₂)₃;
R^{2a} and R^{2b} both represent methyl;
E represents -CH₂N(H)C(O)-(1,3-phenylene)-C(O)N(H)CH₂-;
the -NH- group in the structural fragment of formula Ia is attached in the 6-position of the coumarin ring system;
- 15 R⁷ represent saturated, optionally branched C₁₋₃ alkyl (e.g. methyl, ethyl or isopropyl);
R⁸ represents isopropyl or, particularly, H;
R⁸ represents methyl;
the -NH- and -C(O)- groups in the structural fragment of formula Ic are
20 attached in the 2- and 5-positions, respectively, of the furan or thiophene ring;
R⁹, R¹⁰ and R¹¹ independently represent saturated, optionally branched C₁₋₃ alkyl (e.g. isopropyl);
p represents 1;
- 25 q and r represent 0;
the -C(O)- group in the structural fragment of formula Id is attached in the 4-position of the oxazole ring;
n represents 3 or, when R¹ represents methyl, n represents 2 or 3, or, when R¹ represents D-A-N(H)-[Q]_n-C(O)-E-C(O)-, then n represents 2;


the compound comprises at least one structural fragment of formula Ib, Ic, Id, Ie or If in which R^7 , R^8 , R^9 , R^{10} or R^{11} , respectively, represents isopropyl.

- 5 Particularly preferred compounds of formula I include those in which:
 each Q independently represents a structural fragment of formula Ib, Id, Ie or If (such as a structural fragment of formula Ib or Ie);
 G represents CH;
 R^7 represents methyl or isopropyl.

10

Compounds of formula I that may be mentioned include those in which each Q does not represent a structural fragment of formula Id.

- According to a further aspect of the invention, there is provided compounds
 15 of formula I which are compounds of formula II,



wherein

R^1 represents Het^1 , $R^{1a}\text{C(O)-}$ or $\text{D-A-N(H)-Q}^3\text{-Q}^2\text{-Q}^1\text{-C(O)-E-C(O)-}$;

- 20 Q^1 is absent or represents a structural fragment of formula Ia, Ib, Ic, Id, Ie or If;

Q^2 represents a structural fragment of formula Ib, Ie or If;

Q^3 represents a structural fragment of formula Ib, Id, Ie or If; and

Het^1 , R^{1a} , D, A, E, R^{2a} , R^{2b} , A and the structural fragments of formulae Ia, Ib, Ic, Id, Ie and If are as hereinbefore defined;

25

provided that:

- (a) at least one of Q^1 , Q^2 and Q^3 represents a structural fragment of formula Id, Ie or If; and

- (b) at least one of R^6 or R^7 , R^8 , R^9 , R^{10} and R^{11} (whichever is/are present) represents branched, cyclic or part cyclic C_{3-5} alkyl.

or a pharmaceutically acceptable derivative thereof,

5

which compounds are also referred to hereinafter as "the compounds of the invention".

Compounds of formula II that may be mentioned include those in which at
10 least one of R^7 , R^8 , R^9 , R^{10} and R^{11} (whichever is/are present) represents branched, cyclic or part cyclic C_{3-5} alkyl.

Preferred compounds of formula II include those in which:

Q^1 represents a structural fragment of formula Ic, or, particularly, a
15 structural fragment of formula Ib, Ie or If, or, when R^1 represents methyl or D-A-N(H)- Q^3 - Q^2 - Q^1 -C(O)-E-C(O)-, then Q^1 may also be absent;

Q^2 represents a structural fragment of formula Ib in which G represents CH,
or a structural fragment of formula Ie in which q represents 0;

Q^3 represents a structural fragment of formula Id, Ie or If;
20 at least one of R^7 , R^8 , R^9 , R^{10} and R^{11} (whichever is/are present) represents branched C_{3-5} alkyl.

More preferred compounds of formula II include those in which:

Q^1 represents a structural fragment of formula Ie or, particularly, a
25 structural fragment of formula Ib in which G represents CH, or, when R^1 represents methyl, then Q^1 may also be absent, or, when R^1 represents D-A-N(H)- Q^3 - Q^2 - Q^1 -C(O)-E-C(O)-, then Q^1 is absent;

Q^3 represents a structural fragment of formula Ie.

at least one of R^7 , R^8 , R^9 , R^{10} and R^{11} (whichever is/are present) represents
30 isopropyl or isopentyl;

at least one of R^7 , R^8 , R^9 , R^{10} and R^{11} (whichever is/are present) represents ethyl or, particularly, methyl.

Other preferred compounds of formula II include those in which:

- 5 Q^1 and Q^3 represent a structural fragment of formula Ib in which R^6 represents H and R^7 represents methyl; and
 Q^2 represents a structural fragment of formula Ib in which R^6 represents H and R^7 represents branched, cyclic or part cyclic C_{3-5} alkyl (e.g. cyclopropylmethyl, cyclopentyl or, particularly, isopentyl or isopropyl).

10

In one embodiment of the invention, the compound of formula I or formula II may be one which comprises:

- (a) at least one structural fragment of formula Ib in which G represents N and R^6 represents branched, cyclic or part cyclic C_{3-5} alkyl (e.g. cyclopropylmethyl, isopentyl, cyclopentyl or, particularly, isopropyl);
15 (b) at least one structural fragment of formula Id in which p represents 0 and R^9 represents branched, cyclic or part cyclic C_{3-5} alkyl (e.g. cyclopropylmethyl, isopentyl, cyclopentyl or, particularly, isopropyl); and/or, particularly
20 (c) at least one structural fragment of formula Ie in which q represents 0 and R^{10} represents branched, cyclic or part cyclic C_{3-5} alkyl (e.g. cyclopropylmethyl, isopentyl, cyclopentyl or, particularly, isopropyl).

In this embodiment of the invention, the compound of formula I or formula
25 II may particularly be one which comprises at least one structural fragment of formula Ie in which q represents 0 and R^{10} represents isopropyl.

Compounds of formulae I and II that may be mentioned include those in which p, q and r, in whichever of structural fragments Id, Ie and If,
30 respectively, are present, represent 0.

Preferred compounds of formulae I and II include those that bind to the minor groove of DNA.

Preferred compounds of the invention include those that have different
5 binding affinities at different minor groove binding sites in double-stranded DNA molecules having more than one minor groove binding site.

When used herein, the term “different minor groove binding sites” includes references to minor grooves that differ with respect to sequences of base
10 pairs that they contain (where the sequences may comprise AT base pairs, GC base pairs or mixtures thereof). However, the term also includes references to minor grooves that, although they may contain the same sequence of base pairs, or very similar sequences (e.g. sequences, such as those comprising solely AT base pairs, differing only with respect to the
15 order of the base pairs), they differ with respect to:

- (a) the position of the sequence in the minor groove;
- (b) the sequences of base pairs that are at one or both ends of the sequence correlating to the binding site;
- (c) the pitch angle of base pairs in the minor groove;
- 20 (d) the width of the minor groove.

Compounds of the invention that may be mentioned include those that bind to and/or have specificity for DNA sequences that contain at least one GC base pairing.

25

When used herein, the term “bind to DNA sequences that contain at least one GC base pairing” refers to compounds that, although they bind to DNA minor grooves that contain AT base pairs only, they also bind to DNA minor grooves that contain at least one GC base pairing (i.e. the binding of

the compounds to DNA is “tolerant” of the replacement of AT base pairs with GC base pairs).

When used herein, the term “specificity for DNA sequences that contain at least one GC base pairing” refers to compounds that bind with a measurably greater affinity to DNA minor grooves that contain at least one GC base pairing than they do to DNA minor grooves that contain AT base pairs only.

In relation to the above, affinity to DNA may be measured by techniques known to those skilled in the art, such as capillary electrophoresis, for example as described hereinafter. Furthermore, affinity to certain sections of DNA may be determined by techniques known to those skilled in the art, such as DNA footprinting, for example as described hereinafter.

In this respect, preferred compounds of the invention include:

- (i) compounds of formula I, as hereinbefore defined, provided that the compound comprises at least one structural fragment of formula Id, Ie or If;
- (ii) compounds of formula II, as hereinbefore defined.


Preferred compounds of the invention include the compounds of the Examples disclosed hereinafter.

Preparation


Compounds of the invention may be prepared using techniques (e.g. peptide synthesis) known to those skilled in the art, using starting materials that are either commercially available, are known in the literature, or may be obtained either by analogy with the processes described hereinafter, or by conventional synthetic procedures, in accordance with standard techniques; and appropriate reagents and reaction conditions.

According to the invention there is also provided a process for the preparation of compounds of formula I which comprises:

(a) reaction of a compound of formula III,




- 5 wherein A^a represents A or, when a represents 0, then A^a may also represent A^2 and Q, D, A and A^2 are as hereinbefore defined and a is as defined below, with a compound of formula IV,



- wherein A^b represents a direct bond or $-\text{A}^1-\text{C}(\text{O})-$, as appropriate, L^1 represents a suitable leaving group (e.g. halo, such as chloro, C_{1-6} alkoxy, such as ethoxy, trichloromethyl, hydroxy or an alkoxy group (e.g. a benzyl alkoxy group) that is a functional group of a functionalised polymer (e.g. BALa-PAM resin, or derivatives thereof)), a and b both represent integers from 0 to 5, the sum of the two being 2, 3, 4 or 5, and R^1 and Q are as hereinbefore defined, for example under conditions known to those skilled in the art (for example: (i) when L^1 represents ethoxy, by reaction at between room and reflux temperature in the presence of ethanol; (ii) when L^1 represents halo, by reaction at between -10°C and room temperature in the presence of a suitable base such as diisopropylamine and an appropriate solvent such as dichloromethane; (iii) when L^1 represents hydroxy, by reaction at around room temperature in the presence of an appropriate coupling agent such as HBTU (or a combination of a carbodiimide, such as diisopropylcarbodiimide, and HOBT), a suitable solvent such as DMF, and optionally in the presence of a tertiary amine base such as NMM; and (iv) when L^1 represents a benzyl alkoxy group derived from a BALa-PAM resin, by reaction at between room and reflux

temperature (e.g. 55°C) in the presence of a suitable solvent, such as THF);
or

- (b) for compounds of formula I in which R¹ represents D-A-N(H)-[Q]_n-
5 C(O)-E-C(O)-, reaction of two equivalents of a compound of formula V,



wherein Q, n, A and D are as hereinbefore defined, with a compound of
formula VI,



- 10 wherein L² represents a suitable leaving group (e.g. halo or, particularly, hydroxy), the two L² groups being the same or different, and E is as hereinbefore defined, for example under conditions known to those skilled in the art (e.g. as described in respect of process (a) above).

- 15 Compounds of formulae III, IV, V and VI, and derivatives thereof, are either commercially available, are known in the literature, or may be obtained either by analogy with the processes described herein, or by conventional synthetic procedures, in accordance with standard techniques, from readily available starting materials using appropriate reagents and
20 reaction conditions.

- Substituents on the aryl (e.g. phenyl), and heterocyclic, group(s) in compounds defined herein may be converted to other claimed substituents using techniques well known to those skilled in the art. For example, hydroxy
25 may be converted to alkoxy, phenyl may be halogenated to give halophenyl, nitro may be reduced to give amino, halo may be displaced by cyano, etc.

The skilled person will also appreciate that various standard substituent or functional group interconversions and transformations within certain compounds of formula I will provide other compounds of formula I. For example, alkenylene may be reduced to alkylene, carbonyl may be reduced to hydroxy or methylene, etc.

The compounds of the invention may be isolated from their reaction mixtures using conventional techniques.

It will be appreciated by those skilled in the art that, in the process described above, the functional groups of intermediate compounds may be, or may need to be, protected by protecting groups.

Functional groups which it is desirable to protect include hydroxy, amino and carboxylic acid. Suitable protecting groups for hydroxy include trialkylsilyl and diarylalkylsilyl groups (e.g. *tert*-butyldimethylsilyl, *tert*-butyldiphenylsilyl or trimethylsilyl), tetrahydropyranyl and alkylcarbonyl groups (e.g. methyl- and ethylcarbonyl groups). Suitable protecting groups for amino include benzyl, *tert*-butyloxycarbonyl, 9-fluorenylmethoxycarbonyl or benzyloxycarbonyl. Suitable protecting groups for carboxylic acid include C₁₋₆ alkyl or benzyl esters.

The protection and deprotection of functional groups may take place before or after any of the reaction steps described hereinbefore.

Protecting groups may be removed in accordance with techniques which are well known to those skilled in the art and as described hereinafter.

The use of protecting groups is fully described in "Protective Groups in Organic Chemistry", edited by J.W.F. McOmie, Plenum Press (1973), and

“Protective Groups in Organic Synthesis”, 3rd edition, T.W. Greene & P.G.M. Wutz, Wiley-Interscience (1999).

Persons skilled in the art will appreciate that, in order to obtain compounds of the invention in an alternative, and, on some occasions, more convenient, manner, the individual process steps mentioned herein may be performed in a different order, and/or the individual reactions may be performed at a different stage in the overall route (i.e. substituents may be added to and/or chemical transformations performed upon, different intermediates to those associated hereinbefore with a particular reaction). This will depend *inter alia* on factors such as the nature of other functional groups present in a particular substrate, the availability of key intermediates and the protecting group strategy (if any) to be adopted. Clearly, the type of chemistry involved will influence the choice of reagent that is used in the said synthetic steps, the need, and type, of protecting groups that are employed, and the sequence for accomplishing the synthesis.

Some of the intermediates referred to hereinbefore are novel. According to a further aspect of the invention there is thus provided a compound of formula V, or a protected derivative thereof.

Uses and Pharmaceutical Preparations

Compounds of the invention are useful because they possess pharmacological activity. They are therefore indicated as pharmaceuticals.

Thus, according to a further aspect of the invention there is provided the compounds of the invention for use as pharmaceuticals.

In particular, compounds of the invention bind to DNA, thereby inhibiting the binding to that DNA of an enzyme necessary for replication, with the effect that DNA replication is inhibited.

- 5 Thus, according to a further aspect of the invention there is provided a method of inhibiting DNA replication, which method comprises contacting the DNA with an inhibitory amount of a compound of the invention.

Due to their ability to inhibit DNA replication (e.g. by inhibiting
10 transcription by blocking the binding or displacement of regulatory proteins or DNA-enzyme complexes, such as with reverse transcriptase or topoisomerases), compounds of the invention have utility in the treatment of diseases that rely upon DNA replication for their propagation. Such diseases include cancers and those involving viruses, bacteria, fungi or other
15 microorganisms (e.g. diseases involving parasites, such as malaria).

Thus, according to a further aspect of the invention, there is provided a method of treatment of a disease that relies upon DNA replication for its propagation (e.g. cancer or a viral, bacterial, fungal or other microbial
20 infection), which method comprises administration of a therapeutically effective amount of a compound of the invention to a person suffering from that disease. Such treatment may be particularly useful where the person suffering from that disease is immunocompromised.

- 25 Because they have a different mode of action to many conventional anti-viral, anti-bacterial, anti-fungal or other anti-microbial (e.g. anti-parasitic) agents, compounds of the invention may be particularly useful in the treatment of viral, bacterial, fungal or other microbial (e.g. parasitic) infections where the infective agent is resistant to one or more anti-viral,
30 anti-bacterial, anti-fungal or other anti-microbial (e.g. anti-parasitic) agents

having a different mode of action. In this respect, according to a further aspect of the invention there is provided a method of treating a viral, bacterial, fungal or other microbial (e.g. parasitic) infection, where the viral, bacterial, fungal or other microbial (e.g. parasitic) infective agent is resistant to one or more anti-viral, anti-bacterial, anti-fungal or other anti-microbial (e.g. anti-parasitic) agents, respectively, that do not act by inhibiting DNA replication, which method comprises administration of a therapeutically effective amount of a compound of the invention to a person having that infection.

As well as having utility on their own in the treatment of diseases that rely upon DNA replication for their propagation, the compounds of the invention may be used in combination with one or more other compounds or treatment regimes that are used to treat such a disease. Thus, according to a further aspect of the invention, there is provided a method of treatment of a disease that relies upon DNA replication for its propagation (e.g. cancer or a viral, bacterial, fungal or other microbial infection), which method comprises administration, to a person suffering from that disease, of a therapeutically effective amount of a compound of the invention in combination with one or more other agents that are known to be effective in treating that disease.

When used herein, the term "in combination with" includes administration of the other agent(s) that is(are) known to be effective in treating the disease, before, during and/or following administration a compound of the invention. When more than one other agent is administered, the term also includes administration of the different other agents at different times relative to the time of administration of a compound of the invention.

Agents that are known to be effective in treating diseases that rely upon DNA replication for their propagation (e.g. anti-cancer, anti-viral, anti-bacterial, anti-fungal or other anti-microbial (e.g. anti-parasitic) agents) include those listed under the relevant headings in "*Martindale: The Complete Drug Reference*", 32nd Edition, the Pharmaceutical Press, London (1999), the disclosures of which document are hereby incorporated by reference.

Anti-cancer agents also include non-chemical agents such as ionising radiation (e.g. subatomic particle radiation such as α -particles, β -particles, neutrons, protons, mesons and heavy ions or electromagnetic radiation such as high-frequency X-rays or gamma rays). Chemical anti-cancer agents that may be mentioned include:

(a) Alkylating agents including:

- (i) nitrogen mustards such as mechlorethamine (HN_2), cyclophosphamide, ifosfamide, melphalan (L-sarcolysin) and chlorambucil;
- (ii) ethylenimines and methylmelamines such as hexamethylmelamine, thiotepa;
- (iii) alkyl sulfonates and thiosulfonates such as busulfan, methyl methanesulfonate (MMS) and methyl methanethiosulfonate;
- (iv) nitrosoureas and nitrosoguanidines such as carmustine (BCNU), lomustine (CCNU), semustine (methyl-CCNU), streptozocin (streptozotocin) and *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine (MNNG); and
- (v) triazenes such as dacarbazine (DTIC; dimethyltriazenoimidazole-carboxamide).

(b) Antimetabolites including:

- (i) folic acid analogues such as methotrexate (amethopterin);

- (ii) pyrimidine analogues such as fluorouracil (5-fluorouracil; 5-FU), floxuridine (fluorodeoxyuridine; FUdR) and cytarabine (cytosine arabinoside); and
 - (iii) purine analogues and related inhibitors such as mercaptopurine (6-mercaptapurine; 6-MP), thioguanine (6-thioguanine; TG) and pentostatin (2'-deoxycofomycin).
- (c) Natural Products including:
- (i) vinca alkaloids such as vinblastine (VLB) and vincristine;
 - (ii) epipodophyllotoxins such as etoposide and teniposide;
 - (iii) antibiotics such as dactinomycin (actinomycin A, C, D or F), daunorubicin (daunomycin; rubidomycin), doxorubicin, bleomycin, plicamycin (mithramycin) and mitomycin (mitomycin A, B or C);
 - (iv) enzymes such as L-asparaginase; and
 - (v) biological response modifiers such as interferon alphenomes.
- (d) Miscellaneous agents including:
- (i) platinum coordination complexes such as cisplatin (*cis*-DDP) and carboplatin;
 - (ii) anthracenedione such as mitoxantrone and anthracycline;
 - (iii) substituted urea such as hydroxyurea;
 - (iv) methyl hydrazine derivatives such as procarbazine (N-methylhydrazine, MIH);
 - (v) adrenocortical suppressants such as mitotane (*o,p'*-DDD) and aminoglutethimide;
 - (vi) taxol and analogues/derivatives;
 - (vii) hormone agonists/antagonists such as flutamide and tamoxifen;
 - (viii) photoactivatable compounds (e.g. psoralens);
 - (ix) DNA topoisomerase inhibitors (e.g. m-amsacrine and camptothecin);

- (x) anti-angiogenesis agents (e.g. SU6668, SU5416, combretastatin A4, angiostatin and endostatin); and
- (xi) immunotherapeutic agents (e.g. radiolabelled antibodies such as Bexxar™ and Theragyn™ (Pemtumomab™)).

5

Anti-viral agents that may be mentioned include acyclovir, gancyclovir, AZT, ddI, amantadine hydrochloride, inosine pranobex, vidarabine, and the like.

Anti-bacterial agents that may be mentioned include natural and synthetic
10 penicillins and cephalosporins, sulphonamides, erythromycin, kanomycin, tetracycline, chloramphenicol, rifampicin and including gentamicin, ampicillin, benzypenicillin, benethamine penicillin, benzathine penicillin, phenethicillin, phenoxy-methyl penicillin, procaine penicillin, cloxacillin, flucloxacillin, methicillin sodium, amoxicillin, bacampicillin hydrochloride,
15 ciclacillin, mezlocillin, pivampicillin, talampicillin hydrochloride, carfecillin sodium, piperacillin, ticarcillin, mecillinam, pirmecillinan, cefaclor, cefadroxil, cefotaxime, cefoxitin, cefsulodin sodium, ceftazidime, ceftizoxime, cefuroxime, cephalixin, cephalothin, cephamandole, cephalolin, cephradine, latamoxef disodium, aztreonam, chlortetracycline hydrochloride,
20 clomocycline sodium, demeclocycline hydrochloride, doxycycline, lymecycline, minocycline, oxytetracycline, amikacin, framycetin sulphate, neomycin sulphate, netilmicin, tobramycin, colistin, sodium fusidate, polymyxin B sulphate, spectinomycin, vancomycin, calcium sulphaloxate, sulfametopyrazine, sulphadiazine, sulphadimidine, sulphaguanidine,
25 sulphaurea, capreomycin, metronidazole, tinidazole, cinoxacin, ciprofloxacin, nitrofurantoin, hexamine, streptomycin, carbenicillin, colistimethate, polymyxin B, furazolidone, nalidixic acid, trimethoprim-sulfamethoxazole, clindamycin, lincomycin, cycloserine, isoniazid, ethambutol, ethionamide, pyrazinamide and the like.

30

Anti-fungal agents that may be mentioned include miconazole, ketoconazole, itraconazole, fluconazole, fusidic acid, amphotericin, flucytosine, griseofulvin, natamycin, nystatin, and the like.

5 Anti-parasitic agents (e.g. anti-malarial agents) that may be mentioned include pyrimethamine, proguanil, chloroquine, primaquine, mefloquine, quinine, tetracycline, atovaquone, artemisinin, dihydroartemisinin, artemether, arteether, artesunic acid and its salts, and sulfonamides.

10 When a compound of the invention is administered to a patient in combination with one or more other agents that are known to be effective in treating diseases that rely upon DNA replication for their propagation, the compound of the invention and the other agent(s) may be administered separately or, conveniently, as a single composition. Thus, according to a
15 further aspect of the invention, there is provided a combination product comprising components:

- (A) a formulation comprising a compound of the invention; and
- (B) a formulation comprising one or more other chemical agents that are known to be effective in treating diseases that rely upon DNA
20 replication for their propagation.

The combination product according to this aspect of the invention provides for the administration of a compound of the invention in conjunction with one or more other chemical agents that are known to be effective in treating
25 diseases that rely upon DNA replication for their propagation, and may thus be presented either as separate components (i.e. (A) and (B) separately), or may be presented (i.e. formulated) as a combined preparation (i.e. presented as a single formulation including a compound of the invention and one or more other chemical agents that are known to be effective in treating diseases
30 that rely upon DNA replication for their propagation). When components (A)

and (B) are presented as separate components, the combination product may alternatively be termed "a kit-of-parts".

In this aspect of the invention, other chemical agents that are known to be effective in treating diseases that rely upon DNA replication for their propagation include those referred to or mentioned hereinbefore. Thus, in a preferred embodiment of this aspect of the invention, the other chemical agents that are known to be effective in treating diseases that rely upon DNA replication for their propagation are one or more chemical anti-cancer, anti-viral, anti-bacterial, anti-fungal and/or anti-parasitic agents (e.g. the agents referred to or mentioned hereinbefore).

In a further preferred embodiment of this aspect of the invention, each of components (A) and (B) is formulated in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier.

The compounds of the invention will normally be administered orally, subcutaneously, intravenously, intraarterially, transdermally, intranasally, by inhalation, or by any other parenteral route, in the form of pharmaceutical preparations comprising the active ingredient either as a free base or a non-toxic organic or inorganic acid addition salt, in a pharmaceutically acceptable dosage form. Depending upon the disorder and patient to be treated, as well as the route of administration, the compositions may be administered at varying doses.

According to a further aspect of the invention there is thus provided a pharmaceutical formulation including a compound of the invention in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier. Such formulations may be used for the treatment of diseases that rely upon DNA replication for their propagation. Thus, in one embodiment of this

aspect of the invention there is provided a pharmaceutical formulation including, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier, a compound of the invention and one or more other chemical agents that are known to be effective in treating diseases that rely upon DNA replication for their propagation (e.g. one or more chemical anti-cancer, anti-
5 viral, anti-bacterial, anti-fungal and/or anti-parasitic agents, such as the agents referred to or mentioned hereinbefore).

Suitable daily doses of the compounds of the invention in therapeutic
10 treatment of humans are about 1 to 2000 mg/m².

The most effective mode of administration and dosage regimen for the compounds of the invention depends on several factors, including the particular condition being treated, the extent and localisation of that condition
15 in the patient being treated, as well as the patient's state of health and their reaction to the compound being administered. Accordingly, the dosages of the compounds of the invention should be adjusted to suit the individual patient. Methods for determining the appropriate dose for an individual patient will be known to those skilled in the art.

20

As well as having utility in the treatment of diseases, compounds of the invention are also useful in various assay methods based upon DNA binding. For example, it is known that compounds that bind to the minor groove of DNA have the ability to stabilise DNA duplexes, as well as to
25 stabilise a fully matched (in terms of base pairs) DNA duplex to a greater extent than a mismatched DNA duplex, thereby enabling easier discrimination between the fully matched and mismatched duplexes (e.g. in terms of the melting temperatures of the duplexes).

Thus, according to a further aspect of the invention, there is provided a method of stabilising a DNA duplex formed between first and second single strands of DNA, which method comprises contacting that DNA duplex with a compound of the invention.

5

Further, there is also provided a method of enhancing the difference in melting temperatures between first and second DNA duplexes, wherein each DNA duplex is formed from a first single strand of DNA that is the same in each duplex and a second single strand of DNA that is different in each duplex, which method comprises contacting each DNA duplex with a compound of the invention. In a preferred embodiment, the first DNA duplex has a greater degree of base-pair matching (e.g. it is fully matched) than the second DNA duplex, which has at least one base-pair mismatch.

10

Compounds that stabilise fully matched DNA duplexes to a greater extent than mismatched DNA duplexes may be used to reduce levels of “false positive” results in DNA hybridisation assay techniques, for example as described in US 6,221,589, the disclosures of which are hereby incorporated by reference. The reduction in “false positive” results may be achieved through the use of more stringent conditions (e.g. higher wash temperatures) following a hybridisation reaction in the presence of a duplex-stabilising compound than is possible following a reaction in the absence of such a compound. Thus, there is further provided a method of increasing the maximum temperature of a wash following a DNA hybridisation reaction, the method comprising the provision of a compound of the invention to the hybridisation reaction mixture. When used herein, the term “maximum temperature of a wash following a DNA hybridisation reaction” refers to the highest possible wash temperature that does not result in a substantial loss of the “true positive” results (i.e. the fully or most highly matched DNA duplexes).

25

30

When used herein in relation to the above-mentioned methods involving DNA duplexes, the term "contacting" includes admixing of a compound of the invention with a DNA duplex. However, the term also includes attaching (e.g. covalently bonding) a compound of the invention (e.g. a compound of the invention bearing a haloalkyl group), or a derivative thereof (e.g. a compound of formula V) that bears a functional group (e.g. a hydroxy, amino or carboxylic acid group) that may be used to form a suitable attachment, to one or both of the single strands of DNA that form the duplex. Such "labelled" single strands of DNA may be used as primers, capture probes, or in a number of different assays (e.g. capture-detection assays, 5'-nuclease assays and Beacon assays).

Compounds of the invention have the advantage that, compared to corresponding compounds that, in place of the essential branched, cyclic or part cyclic C₃₋₅ alkyl groups of the compounds of the invention, bear methyl substituents (for example compounds known in the prior art), they have a greater affinity and/or selectivity for any given DNA sequence. For example, compared to compounds known in the prior art, compounds of the invention have the advantage that they bind to DNA minor grooves that contain certain DNA sequences (e.g. sequences comprising AT base pairs only) with a measurably greater affinity and/or selectivity.

Compounds of the invention have the advantage in that they can bind to DNA sequences containing AT base pairs only, GC base pairs only, or mixtures of AT and GC base pairs. However, for pharmaceutical applications compounds of the invention (e.g. compounds of formula I comprising at least one structural fragment of formula Id, If or, particularly, Ie, or compounds of formula II) have the further advantage that, compared to corresponding compounds known in the prior art, they bind to DNA

minor grooves that contain at least one GC base pairing with a measurably greater affinity and/or selectivity.

Compounds of the invention may also have the advantage that they may be more efficacious than, be less toxic than, have a broader range of activity than, be more potent than, be longer acting than, produce fewer side effects than, be more easily absorbed than, or that they may have other useful pharmacological properties over, compounds known in the prior art.

10 Description of the Figures

Figures 1 to 6

Capillary electrophoresis was used to study of ligand-DNA binding (see: (i) Hamdan, I.I.; Skellern, G.G.; and Waigh, R.D. "Separation of pd(GC)₁₂ from pd(AT)₁₂ by free solution capillary electrophoresis", *J. Chromatogr. A.* 806, 165-168 (1998); (ii) Hamdan, I.I.; Skellern, G.G.; and Waigh, R.D. "Use of capillary electrophoresis in the study of ligand-DNA interactions", *Nucleic Acids Research* 26, 3053-3058 (1998); (iii) Hamdan, I.I.; Skellern, G.G.; and Waigh, R.D. "Ligand binding to oligonucleotides" *Methods in Molecular Biology*, Vol. 163, "Capillary Electrophoresis of Nucleic acids", Vol 2, pages 379-391, K.R. Mitchelson and J. Cheng Eds., Humana Press, New Jersey, (2000)).

Typically, ligand binding results in a change in migration time of the DNA and sometimes a change in peak shape.

The following are shown in Figures 1-6:

- (a) the experimental conditions, including the DNA sequence(s);
- (b) the result of sequential addition of ligand; and
- 30 (c) the structure of the ligand being investigated.

Key to Figures 1 to 6:

r0 = electropherogram in the absence of drug

rX = (where X is 0.5, 1, 2, 3 or 4) electropherogram in the presence
of drug, where the drug : DNA ratio (in terms of
concentration) is X:1

As can be seen from the figures, the disappearance of a peak corresponding to uncomplexed DNA relates to a measured ratio (rX) of drug to DNA. This ratio corresponds to the binding ratio.

10 Figure 1: Binding studies with netropsin

Double-stranded DNA oligonucleotides having, on one strand, nucleotide sequences AAATTATATTAT (peak 1) and GGGCCGCGCCGC (peak 2) are incubated in the presence of netropsin, a well-documented AT-selective binder. At r0, the two strands are present without any of the drug. At r1, the presence of netropsin (20 μ M) causes a new peak (the complex of netropsin and the AT dodecamer) to appear in the trace. Increasing the amount of netropsin in the mixture to 40 μ M (r2) causes complete conversion of the free AT dodecamer to its complex with netropsin.

20 From the traces shown in Figure 1, it can be determined that:

- (i) netropsin is AT selective; and
- (ii) it has a binding ratio of 2:1 (drug : DNA) with AAATTATATTAT (1 drug molecule to each available binding site).

25 Figure 2: Binding studies with 12/41 (the compound of Example 2)

Compound 12/41 shows, with AAATTATATTAT, the same 2:1 binding ratio as netropsin (as well as similar AT-selectivity).

Figure 3: Binding studies with 13/20 (the compound of Example 3)

Figure 3 presents the binding electropherograms of 13/20 in the presence of a double-stranded DNA decamer having, on one strand, the nucleotide sequence CGACTAGTCG, the central part of which was indicated to be a high affinity binding site from footprinting studies (see Figure 7). Here we see that the binding ratio is 4:1 (drug : DNA). This result demonstrates that 13/20 has a strong affinity for the given sequence, and also that the minor groove produced by the DNA decamer accommodates 4 drug-binding molecules in one binding site.

Figure 4: Binding studies with distamycin

Figure 4 presents the binding electropherograms of the well-known compound distamycin in the presence of the double-stranded DNA decamer having, on one strand, the nucleotide sequence CGACTAGTCG. It can clearly be seen, distamycin has no affinity for the CGACTAGTCG sequence even at a drug to DNA ratio of 4:1 (the trace labelled r4).

Figure 5: Competitive binding studies with compound 13/20

Figure 5 presents the binding electropherograms for 13/20 in the presence of a mixture of double-stranded DNA oligonucleotides, one having, on one strand, the dodecameric nucleotide sequence AAATTATATTAT (peak 1) and the other having, on one strand, the decameric nucleotide sequence GGACTAGTCG (peak 2). It can be seen from the results that 13/20 has a greater affinity for the decamer ACTA sequence (as indicated by the earlier disappearance of the peak corresponding to the uncomplexed ACTA sequence).

Figure 6: Further binding studies with compound 13/20

Figure 6 presents the binding electropherograms of 13/20 incubated against the double-stranded DNA oligonucleotide having, on one strand, the

decameric nucleotide sequence CCACTAGTGG. The result obtained for this binding was identical with that achieved with the sequence in Fig. 3, showing that it is the central sequence which dominates binding, as indicated by footprinting.

5

Figures 7 and 8: DNA footprinting studies

DNA footprinting is a well-established technique in which a section of radio-labelled DNA is incubated with a chemical or biological cleaving agent, in the presence and absence of a putative binding ligand. Regions of
10 ligand binding will show as blanks in the 'ladder'. Knowing the DNA sequence, the binding sites can then be deduced, from the location of these blanks.

The sequence shown in Figure 7 and 8 is MS1, which contains all possible
15 four-base-pair binding sites. The procedure used in the footprinting experiments was analogous to that described in *J. Med. Chem.* 43, 3257 (2000), the disclosures of which document are hereby incorporated by reference. In an analogous manner to the procedure described in *J. Med. Chem.* 43, 3257 (2000), the MS1 sequence was labelled at the 3'-end with
20 [α - 32 P]dATP using reverse transcriptase. The agent used to cleave the radio-labelled DNA was DNase I.

Bases that fall within the "footprint" of a compound are indicated by xx...xx underneath the sequence. The binding domain that has been confirmed by
25 capillary electrophoresis is indicated by XX...XX.

Figure 7 shows the results for compound 13/20 (at a concentration of 0.03 μ M), whereas Figure 8 shows the results for compound 13/51 (at the concentrations indicated therein). As can be seen, at the concentration used

in the experiment, compound 13/20 produces a footprint that is indicative of binding to only one site on the MS1 sequence. This binding site includes GC base pairs. In contrast, compound 13/51, at the concentrations indicated in Figure 8, produced footprints over several different "AT-only" binding sites on the MS1 sequence.

Figure 9: DNA-binding assays with compounds 12/41 and 13/20.

Murine macrophages (obtained and cultured as described in *Cell. Signal.* 11(7), 491 (1999), hereinafter referred to as "Reference 1", the disclosures of which document are hereby incorporated by reference) were exposed to vehicle (see Reference 1) or 1 µg/mL lipopolysaccharide (LPS; the active component of gram-negative endotoxin and a strong activator of the Nuclear Factor kappa B (NFκB) pathway; the particular LPS being obtained from *Salmonella minnesota*) for 1 hour and nuclear extracts prepared by standard procedures (see *Brit. J. Pharmacol.* 134, 393 (2001), hereinafter referred to as "Reference 2", the disclosures of which document are hereby incorporated by reference).

Aliquots of compounds (1 µL of a 1 mM solution) were incubated with 1 µL (0.035 pmol) of a ³²P-labelled 20bp double-stranded DNA oligonucleotide (representing the consensus binding sequence for NFκB proteins (see Reference 2 and *Brit. J. Pharmacol.* 134, 1629-1638 (2001), the disclosures of which latter document are hereby incorporated by reference)) for 30 min at room temperature. Following this, 5 µg of nuclear protein and buffer (the "gel-shift binding buffer" of Reference 2) were added (final volume 10 µL) and the samples incubated for a further 30 min at room temperature prior to their electrophoresis on pre-run non-denaturing polyacrylamide gels for 45-60 min (see Reference 2). Gels were then dried and exposed to X-ray film. The formation of NFκB protein-DNA

complexes and potential inhibition of complex formation in the presence of the above compounds was detected by autoradiography.

In Figure 9, lanes 1 to 4 show the results obtained where the nuclear protein
5 was extracted from the macrophages exposed to vehicle, whereas lanes 5 to 8 show the results obtained where the nuclear protein was extracted from the macrophages exposed to LPS.

From the results presented in Figure 9, it can be determined that exposure to
10 lipopolysaccharide resulted in an increase in NF κ B–DNA-binding activity (lane 1 vs. lane 5). Pre-incubation with the synthetic compounds had no effect upon basal DNA-binding activity (lane 1 vs. lanes 2 to 4). However, compounds 13/51 and 13/20 appeared to inhibit LPS-stimulated NF κ B-DNA binding activity (lane 5 vs. lanes 6 and 7). In contrast, compound
15 12/41 had little effect (lane 5 vs. lane 8).

Although compound 13/51, which does not bear a branched, cyclic or part
cyclic C₃₋₅ alkyl substituent, appears to inhibit LPS-stimulated NF κ B-DNA
binding activity more strongly than does compound 13/20 (lane 6 vs. lane
20 7), this might well be expected from the lower binding specificity of compound 13/51 (compare Figures 7 and 8).

Biological Tests

25 Minimum inhibitory concentrations (MICs) against microorganisms (e.g. *S. Aureus*, *E. coli*, *S. faecalis*, *P. vulgaris*, MRSA, *Aspergillus niger*, *Candida albicans*, *Klebs. aerogenes*, *Ent. cloacae*, *Mycobacterium fortuitum* or *Aspergillus nidulans*) for compounds of the invention may be measured using procedures such as those described in A. J. Drummond and R. D.

Waigh "The development of microbiological methods for phytochemical screening" *Recent Res. Devel. Phytochem.* 4, 143-152 (2000).

Examples

5

General Experimental Procedures

Electrospray mass spectra (ES-MS) were obtained on a Fisons® VG Platform Benchtop LC-MS. Electron impact (EI-MS) and fast atom bombardment (FAB-MS) mass spectra were obtained on a Jeol® JMS-AX505HA mass spectrometer.

NMR spectra were obtained on a Bruker® AMX 400 spectrometer operating at 400 MHz for ^1H . In ^1H NMR spectra, the abbreviation 'exch.' signifies that the relevant resonance disappeared on treatment of the solution with D_2O .

HPLC purification of the final products was carried out using a Vydac protein and peptide C18 column on a gradient eluting system. The solvents were A: water + 0.1% trifluoroacetic acid (TFA), and B: acetonitrile + 0.1% TFA.

IR spectra: solids were run as KBr discs and liquids as films, using a Nicolet® Impact 400D.

25

Column chromatography was performed with silica gel Prolabo® (200-400 mash).

Example 1

N-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]amino}carbonyl)-1-isopropyl-1-H-pyrrol-3-yl]-4-[(3,3-dimethylbutanoyl)amino]-1-methyl-1H-pyrrole-2-carboxamide

5

(i) Ethyl 4-nitro-1H-pyrrole-2-carboxylate

The sub-title compound was prepared according to a standard literature procedure (see: Ref.1: W.J. Hale and W.V. Hoyr, *J.Amer.Chem.Soc.* 37, 2538-2552 (1915); and Ref.2: K.J. Morgan and D.P. Morrey, *Tetrahedron* 22, 57-62 (1966)).

10

(ii) Ethyl 1-isopropyl-4-nitro-1H-pyrrole-2-carboxylate

Ethyl 4-nitro-1H-pyrrole-2-carboxylate (1.039 g, 5.645 mmol; see step (i) above) was dissolved in DMF (40 mL, dry) to which was added potassium metal (384 mg, 9.831 mmol). The reaction mixture was heated to 100°C and left at that temperature for 1 h with stirring. The reaction mixture was cooled to 50-60°C then isopropyl bromide (1.224 g, 9.953 mmol) and potassium iodide (1.714 g, 10.327 mmol) were added and the reaction mixture was again heated with stirring to 100°C for 5 h, and was then left stirring at room temperature overnight. The solvent was removed under reduced pressure and the residue was diluted with DCM and then extracted with water. The organic extract was dried over MgSO₄, filtered and the solvent removed under reduced pressure. The crude product obtained was chromatographed over silica gel eluted with 1/2 pet.ether / DCM. The material (R_F = 0.26) was collected and the solvent removed under reduced pressure to give white crystalline solid (686 mg, 54% yield), m.p. 60-63°C.

15

20

25

Elemental analysis:

Found: %C=53.19; %H=6.31; %N=12.30;

Calculated for C₁₀H₁₄N₂O₄: %C=53.10; %H=6.24; %N=12.38.

NMR(DMSO- d_6): 7.80(1H, d); 7.45(1H,d); 5.55-5.44(1H, qt); 4.36-4.27(2H, q); 1.51(6H, d); 1.40-1.34(3H, t).

IR ν_{\max} [KBr]: 1714, 1536, 1504, 1440, 1325, 1287, 1223, 1178, 1063 cm^{-1} .

5 (iii) 1-Isopropyl-4-nitro-1H-pyrrole-2-carboxylic acid

Ethyl 1-isopropyl-4-nitro-1H-pyrrole-2-carboxylate (656 mg, 2.900 mmol; see step (ii) above) was dissolved in ethanol (4 mL) to which was added a solution of NaOH (490 mg, 12.250 mmol in water (10 mL)). The reaction mixture was heated under reflux for 1.5 h and then the ethanol was removed
10 under reduced pressure at 40°C. The reaction mixture was extracted with DCM (10 mL), and the water layer was collected, cooled to 0°C with ice-water then acidified with HCl_{conc} with stirring. The white solid material precipitated was filtered off, washed with water and dried under reduced pressure at 50°C to give the required product (548 g, 95% yield), m.p. 194-
15 195°C.

Elemental analysis:

Found: %C=48.40; %H=4.85; %N=14.03;

Calculated for $\text{C}_8\text{H}_{10}\text{N}_2\text{O}_4$: %C=48.48; %H=5.09; %N=14.13.

NMR(DMSO- d_6): 8.37(1H, d); 7.27(1H, d); 5.47-5.36(1H, q); 1.44(6H, d).

20 IR ν_{\max} [KBr]: 1676, 1541, 1517, 1483, 1411, 1370, 1250, 1188, 1077 cm^{-1} .

(iv) N-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]-1-isopropyl-4-nitro-1H-pyrrole-2-carboxamide

N-[3-(Dimethylamino)propyl]-1-methyl-4-nitro-1H-pyrrole-2-carboxamide
25 (816 mg, 3.209 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) was suspended in a mixture of ethanol (30 mL) and HCl (7 mL, dil) to which was added Pd/C (517 mg, 10%) at room temperature. The reaction mixture was hydrogenated at room temperature and atmospheric pressure for 3.5 h. The catalyst was removed by filtration over Kieselguhr and then the solvent
30 was removed under reduced pressure at 50°C to give the amine as pale

yellow solid. The product so formed was dissolved in water (10 mL) to which NaHCO_3 (2 g) was added. 1-Isopropyl-4-nitro-1*H*-pyrrole-2-carboxylic acid (548 mg, 2.631 mmol; see step (iii) above) was dissolved in DME (6 mL, dry) and thionyl chloride (3 mL), then heated under reflux for 5 2 h. The solvent was removed at room temperature to give the acid chloride which was dissolved in benzene (10 mL) and was added dropwise to the reaction mixture at room temperature with stirring. After addition of the acid chloride solution was finished the reaction mixture was heated under reflux for 30 min. then left to cool to room temperature overnight with 10 continuous stirring. The reaction mixture was extracted with DCM (300 mL), dried and the solvent removed under reduced pressure to give yellow solid material. This material was triturated with a small amount of methanol and filtered to give pale yellow solid (318 mg). The filtrate was concentrated and chromatographed over silica gel using ethyl 15 acetate/methanol/ammonia (49/49/2). Fractions with R_f value of 0.3 were collected. The product was obtained as yellow solid (640 mg, 49% yield in total). M.p. 205-207°C.

Found: C,56.25; H,6.78; N,20.59 Calculated for $\text{C}_{19}\text{H}_{28}\text{N}_6\text{O}_4$ C,56.42; H,6.98; N,20.78%.

20 $^1\text{H-NMR}$ (DMSO- d_6) δ 1.42-1.57 (6H,d); 1.60-1.65 (2H,m); 2.12 (6H,s); 2.20-2.25 (2H,t); 3.17-3.22 (2H,q); 3.81 (3H,s); 5.40-5.51 (1H,m); 6.83 (1H,s); 7.22 (1H,s); 7.48 (1H,s); 8.11-8.15 (1H,t, CONH); 8.34 (1H,s); 10.29 (1H,s).

IR ν_{max} [KBr]:1640, 1555, 1506, 1480, 1322, 1279, 1240, 1203 cm^{-1} .

25 ESMS: Found: 405.2 Calculated 405.5.

(v) *N*-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-1-isopropyl-4-{{[(1-methyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]-amino}-1*H*-pyrrole-2-carboxamide

N-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-1-isopropyl-4-nitro-1*H*-pyrrole-2-carboxamide (272 mg, 0.672 mmol; see step (iv) above) and Pd/C (325 mg, 10%) were suspended in isopropanol (30 mL) and hydrogenated at room temperature and atmospheric pressure for 5 h. The catalyst was removed over Kieselguhr and the solvent removed under reduced pressure to give the amine as pale yellow solid, which was used without further purification. The amine was dissolved in THF (5 mL, dry) then 2,2,2-trichloro-1-(1-methyl-4-nitro-1*H*-pyrrol-2-yl)ethanone (183 mg, 0.674 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) dissolved in THF (5 mL, dry) was added dropwise with stirring at room temperature. The stirring was continued overnight at room temperature. The solvent was removed under reduced pressure at room temperature to give the crude product, which was purified by silica gel column chromatography ethyl acetate/methanol/ammonia (49/49/2). Fractions containing the product at R_f value of 0.3 were collected. The product was obtained as yellow solid material (221 mg, 63% yield) with no distinct melting point.

$^1\text{H-NMR}$ (DMSO- d_6) δ 1.36-1.38 (6H,d); 1.64-1.71 (2H,m); 2.31 (6H,s); 3.16-3.22 (2H,q); 3.81 (3H,s); 3.96 (3H,s); 5.40-5.50 (1H,m); 6.85 (1H,d J 1.6Hz); 6.93 (1H,d J 1.6Hz); 7.18 (1H,d J 1.6Hz); 7.43 (1H,d J 1.6Hz); 7.58 (1H,d J 1.6Hz); 8.07-8.10 (1H,t, CONH); 8.18 (1H,d J 1.6Hz); 9.96 (1H,s); 10.29 (1H,s).

IR ν_{max} [KBr]: 1643, 1579, 1534, 1507, 1464, 1407, 1310, 1247 cm^{-1} .

HRFABMS: Found: 527.27341 Calculated for $\text{C}_{25}\text{H}_{35}\text{N}_8\text{O}_5$ 527.27304.

(vi) *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopropyl-1-*H*-pyrrol-3-yl]-4-[(3,3-dimethylbutanoyl)amino]-1-methyl-1*H*-pyrrole-2-carboxamide

N-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-1-isopropyl-4-{{[(1-methyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]-amino}-1*H*-pyrrole-2-carboxamide (100 mg, 0.189 mmol; see step (v) above) was dissolved in methanol (20 mL) to which was added Pd/C (114 mg, 10%). The reaction mixture was hydrogenated at room temperature and atmospheric pressure for 6.5 h. The catalyst was removed over Kieselguhr and the solvent removed under reduced pressure. The amine so formed was dissolved in DCM (10 mL, dry) to which was added a solution of 3,3-dimethylbutanoyl chloride (35 μ L, 0.260 mmol), which was dissolved in DCM (5 mL, dry). The addition was dropwise with stirring at room temperature. Di-isopropylamine was then added in one portion to the reaction mixture. Stirring was continued at room temperature overnight. The solvent was removed under reduced pressure and the crude product obtained was purified by column chromatography using silica gel and ethyl acetate/methanol/ammonia (49/49/2). The solvent was removed under reduced pressure and the solid material was dissolved in chloroform (100 mL) and extracted with (5%, 50 mL) NaHCO₃. The organic layer was dried over Na₂SO₄, filtered and the solvent removed to give the required product (84 mg, 75% yield) as yellow glassy material with no distinct melting point.

¹H-NMR(CDCl₃): δ 1.03 (9H,s); 1.29-1.32 (6H,d); 1.68-1.83 (2H,m); 2.15 (2H,s); 2.19 (6H,s); 2.26-2.39 (2H,t); 3.39-3.46 (2H,q); 3.83 (3H,s); 3.84 (3H,s); 5.42-5.51 (1H,m); 6.47 (1H,d J 1.4Hz); 6.58 (1H,d J 1.4Hz); 6.67 (1H,d J 1.4Hz); 7.05 (1H,d J 1.4Hz); 7.22 (1H,d J 1.4Hz); 7.37 (1H,d J 1.4Hz); 7.78 (1H,t CONH); 7.82 (1H,s); 8.18 (1H,s); 8.32 (1H,s).

IR ν_{\max} [KBr]: 3298, 2960, 1644, 1582, 1538, 1405, 1250 cm⁻¹.

ES-MS: Found: M^+ 595.1 Calculated for $C_{31}H_{46}N_8O_4$ M^+ 595.4.

Example 2

N-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopropyl-1*H*-pyrrol-3-yl]-4-(formylamino)-1-methyl-1*H*-pyrrole-2-carboxamide

N-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-1-isopropyl-4-{{[(1-methyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]-amino}-1*H*-pyrrole-2-carboxamide (102 mg, 0.194 mmol; see Example 1, step (v) above) was dissolved in methanol (20 mL), to which was added Pd/C (106 mg, 10%). The reaction mixture was hydrogenated for 5 h at room temperature and atmospheric pressure. The catalyst was removed over Kieselguhr and then ethyl formate (10 mL) was added to the filtrate. The reaction mixture was then heated until reflux for 72 h. The solvent was removed under reduced pressure and the crude product obtained was purified by HPLC. The product was obtained as white solid material after freeze-drying (64 mg, 51% yield as TFA salt).

$^1\text{H-NMR}$ (DMSO- d_6) δ 1.35-1.37(6H,d); 1.81-1.85(2H,m); 2.78-2.79(6H,d); 3.05(2H,m); 3.23-3.25(2H,q); 3.81(3H,s); 3.84(3H,s); 5.40-5.46(1H,m); 6.92(1H,s); 6.94(1H,s); 6.96(1H,s); 7.17(1H,s); 7.19(1H,s); 7.38(1H,s); 8.12(1H,s); 8.15(1H,t, CONH); 9.3(1H, broad, TFA); 9.85(1H,s); 9.91(1H,s); 10.05(1H,s).

IR ν_{max} [KBr]: 1673, 1644, 1582, 1540, 1470, 1441, 1402, 1204, 1139 cm^{-1} .

ES-MS: Found: M^+ 524.9 Calculated 525.6.

Example 3

N-[3-(Dimethylamino)propyl]-2-({[4-({[4-(formylamino)-1-methyl-1H-pyrrol-2-yl]carbonyl}amino)-1-methyl-1H-pyrrol-2-yl]carbonyl}amino)-5-isopropyl-1,3-thiazole-4-carboxamide

5

(i) Methyl 2-amino-5-isopropyl-1,3-thiazole-4-carboxylate

The sub-title compound was prepared in 41% yield according to the procedure described in *J. Chem. Soc., Perkin Trans. I* 159 (1982).

m.p. 151-152°C (Lit. 150-151°C).

10

(ii) Methyl 5-isopropyl-2-([(1-methyl-4-nitro-1H-pyrrol-2-yl)carbonyl]-amino)-1,3-thiazole-4-carboxylate

1-Methyl-4-nitro-1H-pyrrole-2-carboxylic acid (520 mg, 3.057 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) was dissolved in thionyl chloride (10 mL). The reaction mixture was heated under reflux overnight. Excess thionyl chloride was removed under reduced pressure to give the acid chloride, which was used without further purification. The acid chloride was dissolved in DCM (20 mL, dry), to which NMM (1.214 g, 12.000 mmol). The reaction mixture was cooled to 0°C and kept under N₂.
A solution of methyl 2-amino-5-isopropyl-1,3-thiazole-4-carboxylate (0.612 g, 3.057 mmol; see step (i) above) in DCM (20 mL, dry) was then added dropwise with stirring over a period of 10 min. The temperature was left to rise to room temperature overnight. DCM (10 mL) was added to the reaction mixture then HCl (50 mL, dil.) was added and the reaction mixture was extracted. The organic layer was washed with brine (50 mL), dried over MgSO₄ and the solvent removed under reduced pressure to give the crude product. Column chromatography over flash silica gel [1/1 ethyl acetate / pet. ether] gave the required product as white crystalline solid (460 mg, 43% yield), m.p. 118-120°C (softening).

NMR(CDCI₃): 1.33-1.35(6H, CH-(CH₃)₂, d, J=6.4Hz); 3.89(3H, s, CO₂Me); 4.09(3H, s, NMe); 4.11-4.18(1H, m); 7.24(1H, pyrrole, d, J=1.4Hz); 7.67(1H, pyrrole, d, J=1.4Hz); 9.85(1H, broad, CONH, exch.).

IR ν_{\max} [KBr]: 3557, 3119, 2975, 1712, 1646, 1574, 1509, 1319, 1208 cm⁻¹.

5 HREIMS: Found: 352.08579 calculated for C₁₄H₁₆N₄O₅S 352.08414.

(iii) 5-Isopropyl-2-{[(1-methyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]amino}-1,3-thiazole-4-carboxylic acid

Methyl 5-isopropyl-2-{[(1-methyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]amino}-
10 1,3-thiazole-4-carboxylate (480 mg, 1.363 mmol; see step (ii) above) was
suspended in ethanolic KOH (0.5 M, 25 mL). The reaction mixture was
heated under reflux for 4h, then cooled to 0°C. HCl (conc.) was added
dropwise with stirring until pH 2. The pale yellow solid obtained was
filtered off, washed with de-ionised water and dried under reduced pressure
15 at 45°C overnight to give the required material (440 mg, 96% yield),
m.p.315-319°C[decomp.].

NMR(DMSO-d₆) 1.23-1.25(6H, CH-(CH₃)₂, d, J=6.4Hz); 3.98(3H, s, NMe); 4.01-4.08(1H, m); 7.99(1H, pyrrole, d, J=1.4Hz); 8.29(1H, pyrrole, d, J=1.4Hz); 12.5-13.2(1H, broad, CO₂H, exch.).

20 IR ν_{\max} [KBr]: 3189, 3132, 2972, 1668, 1566, 1537, 1512, 1497, 1319, 1231 cm⁻¹.

HRFABMS: Found: 337.06396 calculated for C₁₃H₁₃O₅N₄S 337.06067

(iv) N-[3-(Dimethylamino)propyl]-5-isopropyl-2-{[(1-methyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]amino}-1,3-thiazole-4-carboxamide

25 5-Isopropyl-2-{[(1-methyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]amino}-1,3-thiazole-4-carboxylic acid (440 mg, 1.301 mmol; see step (iii) above) was
suspended in thionyl chloride (6 mL) and heated until reflux for 3 h.
Solvent (excess thionyl chloride) was removed under reduced pressure and

the acid chloride so formed was dissolved in THF (20 mL, dry) then NMM (330 μ L) was added. A solution of 3-(dimethylamino)propylamine (270 mg, 2.602 mmol) in THF (5 mL, dry) was added dropwise with stirring at room temperature under nitrogen. The stirring was continued at room temperature overnight. The solvent was removed under reduced pressure and the residue was purified by column chromatography using (1:1 methanol / ethyl acetate containing 1%TEA). The product was obtained as yellow glassy material (159 mg, 29% yield), m.p.85-90°C [softening].

NMR(CDCl₃): 1.27-1.29(6H, CH-(CH₃)₂, d, J=6.4Hz); 1.79-1.86(2H, CH₂, qt, J=6.9Hz); 2.31(6H, s, NMe₂); 2.45-2.49(2H, CH₂, t, J=6.9Hz); 3.44-3.50(2H, CH₂, q, J=6.9Hz); 4.10(3H,s, NMe); 4.38-4.45(1H, CH(CH₃)₂, m); 7.56(1H, pyrrole, d, J=1.4Hz); 7.62(1H, s, CONH, exch.); 7.69(1H, pyrrole, d, J=1.4Hz).

IR ν_{\max} [KBr]: 3131, 2959, 1668, 1551, 1500, 1418, 1310, 1286 cm⁻¹.

HREIMS: Found: 422.17695 calculated for C₁₈H₂₆O₄N₆S 422.17363.

(v) *N*-[3-(Dimethylamino)propyl]-5-isopropyl-2-{[(1-methyl-4-[(1-methyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]amino}-1*H*-pyrrol-2-yl)carbonyl]-amino}-1,3-thiazole-4-carboxamide

N-[3-(Dimethylamino)propyl]-5-isopropyl-2-{[(1-methyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]amino}-1,3-thiazole-4-carboxamide (159 mg, 0.376 mmol; see step (iv) above) was dissolved in methanol (25 mL), cooled with ice, and then Pd/C (158 mg, 10%) was added under nitrogen. The reaction mixture was hydrogenated for 4 h at room temperature and atmospheric pressure. The catalyst was removed over Kieselguhr and the solvent was then removed under reduced pressure to give the amine. This amine was dissolved in THF (25 mL, dry) to which was added 2,2,2-trichloro-1-(1-methyl-4-nitro-1*H*-pyrrol-2-yl)ethanone (134 mg, 0.494 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) dissolved in THF

(5 mL, dry), dropwise with stirring, at room temperature. Stirring was continued at room temperature overnight. The solvent was removed under reduced pressure and the residue was chromatographed using 49/49/2 methanol/ethyl acetate/TEA. The product obtained was dissolved in DCM and extracted with a solution of sodium carbonate (10 mL, 5%), then dried and the solvent removed under reduced pressure. The product was obtained as yellow solid (90 mg, 44% yield). M.p.138-141°C.

¹H-NMR(DMSO-d₆) δ 1.26 & 1.28(6H,d,2xMe); 1.65(2H,m,CH₂); 2.19(6H,s,NMe₂); 2.32(2H,t,CH₂); 3.26(2H,q,CH₂); 3.90(3H,s,NMe); 3.97(3H,s,NMe); 4.19(1H,m,CH); 7.36(1H,d,PyrH); 7.47(1H,d,pyrH); 7.62(1H,d,PyrH); 7.81(1H,t,CONH exch.); 8.19(1H,d,PyrH); 10.35(1H,s,CONH exch.); 12.10(1H,broad exch.).

IR ν_{max} [KBr]: 1651,1552,1506,1464,1421,1398,1311,1288,1202 cm⁻¹.

HRFABMS: Found: 545.22931 Calculated for C₂₄H₃₃N₈O₅S 545.22946.

15

(vi) N-[3-(Dimethylamino)propyl]-2-({[4-({[4-(formylamino)-1-methyl-1H-pyrrol-2-yl]carbonyl}amino)-1-methyl-1H-pyrrol-2-yl]carbonyl}amino)-5-isopropyl-1,3-thiazole-4-carboxamide

N-[3-(Dimethylamino)propyl]-5-isopropyl-2-{{[(1-methyl-4-{{[(1-methyl-4-nitro-1H-pyrrol-2-yl)carbonyl]amino}-1H-pyrrol-2-yl)carbonyl]amino}-1,3-thiazole-4-carboxamide (90 mg, 0.165 mmol; see step (v) above) was suspended in ethanol (20 mL) then Pd/C (97 mg, 10%) was added at 0°C under nitrogen. The reaction mixture was hydrogenated at room temperature and atmospheric pressure for 5 h. The catalyst was removed under reduced pressure then ethyl formate (20 mL) was added to the ethanolic solution of the amine. The reaction mixture was heated under reflux for 3 days, then the solvent was removed under reduced pressure. The crude product was dissolved in water (2 mL) containing 0.1% TFA,

then purified by HPLC. The fractions collected were freeze-dried to give the desired product as pale yellow solid (55 mg, 51%) with no distinct m.p.

¹H-NMR(DMSO-d₆) δ 1.27-1.29(6H, d, J=6.9Hz), 1.86(2H, m), 2.78-2.79(6H, d, J=4.0Hz), 3.07(2H, m), 3.37(2H, m), 3.84(3H, s), 3.88(3H, s),
5 4.19(1H, m), 6.94(1H, d, J=1.6Hz), 7.19(1H, s), 7.38(1H, s), 7.40(1H, s),
7.96(1H, t), 8.13(1H, d, J=1.6Hz), 9.30(1H, broad, TFA), 9.99(1H, s),
10.04(1H,s), 12.02(1H, s).

IR ν_{max} [KBr]: 1664, 1555, 1465, 1399, 1294, 1209, 1130 cm⁻¹.

HRFABMS: Found: 544.25962 Calculated for C₂₅H₃₆N₈O₄S 544.25802

10

Example 4

N-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopropyl-1H-pyrrol-3-yl]-4-({[4-(formylamino)-1-isopropyl-1H-pyrrol-2-yl]carbonyl}amino)-1-isopropyl-1H-pyrrole-2-carboxamide

15

(i) N-[3-(Dimethylamino)propyl]-1-isopropyl-4-nitro-1H-pyrrole-2-carboxamide

1-Isopropyl-4-nitro-1H-pyrrole-2-carboxylic acid (312 mg, 1.574 mmol; see Example 1, step (iii) above) was dissolved in thionyl chloride (5 mL) and
20 heated until reflux for 4 h. The excess thionyl chloride was removed under reduced pressure at 50°C to give the acid chloride as white solid material that was used without further purification. 3-(Dimethylamino)propylamine (250 μL) was dissolved in THF (250 mL, dry) to which NMM (250 μL) was added at room temperature with stirring. The acid chloride was
25 dissolved in THF (5 mL, dry) and added dropwise to the amine solution at room temperature with stirring. The reaction mixture was then left stirring at room temperature overnight. Following this the solvent was removed under reduced pressure at 50°C and then the crude product was extracted with K₂CO₃ (25 mL, 10%) and DCM (2X50 mL). The organic layer was

collected, dried over Na₂SO₄, filtered and the solvent removed under reduced pressure. The crude product was purified by chromatography over silica gel using 49.5:49.5:1 methanol / ethyl acetate / TEA to give the product as pale yellow oil, R_f = 0.25, (400 mg, 90% yield).

5 NMR(CDCl₃): 1.46&1.48(6H, d, J=6.7Hz), 1.71-1.77(2H, qt, J=6.0Hz), 2.32(6H, s), 2.51(2H, t, J=5.7Hz), 3.49(2H, qt, J=5.0Hz), 5.62(1H, qt, J=6.7Hz), 6.92(1H, d, J=1.6Hz), 7.73(1H, d, J=1.6Hz), 8.58(1H, s).

IR ν_{max} [KBr]: 3330, 3136, 2979, 1650, 1535, 1430, 1279, 1258 cm⁻¹.

HREIMS: Found: 282.16908 Calculated for C₁₃H₂₂N₄O₃ 282.16919

10

(ii) N-[5-({[3-Dimethylamino]propyl}amino)carbonyl)-1-isopropyl-1H-pyrrol-3-yl]-1-isopropyl-4-nitro-1H-pyrrole-2-carboxamide

N-[3-(Dimethylamino)propyl]-1-isopropyl-4-nitro-1H-pyrrole-2-carboxamide (400 mg, 1.417 mmol; see step (i) above) was dissolved in ethanol (20 mL) then cooled to 0°C. Pd/C-10% (343 mg) was added
15 portion-wise under N₂ with stirring. The reaction mixture was hydrogenated at room temperature and atmospheric pressure for 4 h. The catalyst was removed over Kieselguhr, and the solvent removed under reduced pressure at 50°C to give an oil which was used in the next step
20 without further purification. 1-Isopropyl-4-nitro-1H-pyrrole-2-carboxylic acid (281 mg, 1.417 mmol; see Example 1, step (iii) above) was suspended in thionyl chloride (5 mL) then heated under reflux for 4 h. The excess thionyl chloride was removed under reduced pressure at 50°C and then it was dissolved in THF (5 mL, dry). The amine was dissolved in THF (20
25 mL, dry) to which was added NMM (250 µL) followed by the acid chloride solution. The addition was dropwise at room temperature with stirring. The reaction mixture was left stirring at room temperature overnight. The solvent was removed under reduced pressure and the crude product dissolved in DCM (50 mL) and extracted with Na₂CO₃ (5%, 25 mL). The

organic layer was dried over Na₂SO₄, filtered and the solvent removed under reduced pressure. The crude product was chromatographed over silica gel using 50:50:0.5 methanol / ethyl acetate / TEA. Fractions with R_f value of 0.1 were collected. The solvent was removed at 50°C under reduced pressure to give the product as yellow glassy material (471 mg, 77% yield), m.p.86-90°C (softening).

NMR(CDCl₃):1.42(6H, d, 6.7Hz), 1.75(6H, d, J=6.7Hz), 1.72-1.77(2H, m), 2.31(6H, s), 2.46(2H, t, J=6.1Hz), 3.44-3.49(2H, q, J=6.0Hz), 5.51-5.64(2H, m), 6.45(1H, d, 1.6Hz), 7.17(1H, d, J=1.6Hz), 7.40(1H, d, J=1.6Hz), 7.69(1H, s, broad, CONH), 7.79(1H, d, J=1.6Hz).

IR ν_{max} [KBr]: 2974, 2927, 1636, 1574, 1531, 1507, 1407, 1282, 1237, 1170 cm⁻¹.

HREIMS: Found: 432.24833 Calculated for C₂₁H₃₂N₆O₄ 432.24850

- (iii) *N*-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopropyl-1*H*-pyrrol-3-yl]-1-isopropyl-4-{{[(1-isopropyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]-amino}-1*H*-pyrrole-2-carboxamide
- N*-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopropyl-1*H*-pyrrol-3-yl]-1-isopropyl-4-nitro-1*H*-pyrrole-2-carboxamide (209 mg, 0.483 mmol; see step (ii) above) was dissolved in ethanol (20 mL) at 0°C with stirring. Pd/C-10% (180 mg) was added to the solution at 0°C and under N₂ with stirring. The reaction mixture was hydrogenated for 4 h then the catalyst was removed by filtration over Kieselguhr and the solvent removed under reduced pressure at 50°C to give the amine. 1-Isopropyl-4-nitro-1*H*-pyrrole-2-carboxylic acid (92 mg, 0.462 mmol; see Example 1, step (iii) above) was dissolved in thionyl chloride (3 mL) and heated under reflux for 3 h. Excess thionyl chloride was removed under reduced pressure at 50°C to give the acid chloride. The amine was dissolved in THF (20 mL, dry) to which the acid chloride [dissolved in THF (20 mL, dry)] was added dropwise with stirring at room temperature. The reaction mixture was left

stirring at room temperature overnight. The solvent was removed under reduced pressure and the crude product dissolved in DCM (2x50 mL) after which it was extracted with Na₂CO₃ (50 mL, 5%). The organic layer was dried and then the solvent removed. The crude product was chromatographed over silica gel using 49.5:49.5:1 ethyl acetate/methanol/TEA as eluant. The product was obtained as yellow glassy material $R_f = 0.15$, (230 mg, 82% yield), m.p. 130-133°C (softening). NMR(CDCl₃): 1.43(6H, d, J=6.7Hz), 1.47(6H, d, J=6.7Hz), 1.52(6H, d, J=6.7Hz), 1.73-1.79(2H, m), 2.32(6H, s), 2.47-2.50(2H, t, 6.1Hz), 3.46-3.50(2H, q, 6.0Hz), 5.53-5.61(3H, m), 6.42(1H, d, J=1.6Hz), 6.60(1H, d, J=1.6Hz), 7.28(1H, d, J=1.6Hz), 7.40(1H, d, J=1.6Hz), 7.41(1H, s), 7.44(1H, s), 7.63(1H, s, broad, CONH), 7.81(1H, d, J=1.6Hz), 8.02(1H, s). IR ν_{max} [KBr]: 2979, 2937, 1645, 1593, 1525, 1504, 1410, 1279, 1237, 1190 cm⁻¹.

HRFABMS: Found: 583.33440 Calculated for C₂₉H₄₃N₈O₅ 583.33564

(iv) *N*-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopropyl-1*H*-pyrrol-3-yl]-4-({[4-(formylamino)-1-isopropyl-1*H*-pyrrol-2-yl]carbonyl}-amino)-1-isopropyl-1*H*-pyrrole-2-carboxamide :

N-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopropyl-1*H*-pyrrol-3-yl]-1-isopropyl-4-({[(1-isopropyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]amino}-1*H*-pyrrole-2-carboxamide (105 mg, 0.180 mmol; see step (iii) above) was suspended in ethanol (20 mL) to which Pd/C-10% (80 mg) was added under nitrogen at 0°C. The reaction mixture was hydrogenated at room temperature and atmospheric pressure for 4 h. The catalyst was removed over Kieselguhr and the ethanolic solution was then used in the second part of the experiment. Ethyl formate (20 mL) was added to the ethanolic solution of the amine. The reaction mixture was heated under reflux for 4 days. The solvents were removed under reduced pressure at 40°C and then the crude product was purified by reverse phase HPLC. The fractions

obtained were freeze-dried to give the required material as white solid (71 mg, 57% yield) as TFA salt with no distinct m.p.

NMR(DMSO- d_6): 1.14-1.36(18H, m), 1.82-1.91(2H, m), 2.78(6H, d, $J=4.62\text{Hz}$), 3.04-3.09(2H, m), 3.22-3.26(2H, m), 5.42-5.51(3H, m),
5 6.82(1H, d, $J=1.6\text{Hz}$), 5.89(1H, d, $J=1.6\text{Hz}$), 6.97(1H, d, $J=1.6\text{Hz}$), 7.35(1H, d, $J=1.6\text{Hz}$), 7.34(1H, s), 8.12-8.14(2H, s&t), 9.25(1H, s, broad, TFA), 9.91(1H, s), 9.93(1H, s), 10.03(1H, s).

IR ν_{max} [KBr]: 3285, 2953, 1650, 1580, 1525, 1460, 1402, 1260 cm^{-1} .

HREIMS: Found: 580.35047 Calculated for $\text{C}_{30}\text{H}_{44}\text{O}_4\text{N}_8$ 580.34855.

10

Example 5

N-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]-4-(formylamino)-1-isopentyl-1*H*-pyrrole-2-carboxamide

15

(i) Ethyl 1-isopentyl-4-nitro-1*H*-pyrrole-2-carboxylate

Ethyl 4-nitro-1*H*-pyrrole-2-carboxylate (2.022 g, 10.983 mmol; see Example 1, step (i) above) was dissolved in DMF (30 mL, dry) to which potassium (0.569 g, 14.552 mmol) was added with stirring under nitrogen.

20

The reaction mixture was heated at 100°C for 1 h and then the temperature was allowed to reach 50°C. Isoamyl bromide (2.498 g, 16.538 mmol) was then added, followed by KI (2.197 g, 14.552 mmol). The temperature was raised to 100°C and stirring continued for 5 h, after which time the temperature was allowed to return to room temperature. The reaction

25

mixture was then stirred overnight, following which the solvent was removed under reduced pressure at 50°C. The crude product was dissolved in DCM and the salt extracted with water. The organic extract was dried over MgSO_4 , filtered and the solvent removed under reduced pressure at 30°C. The crude product obtained was chromatographed over silica gel

30

using 1/2 *n*-hexane/DCM as eluant. Material with $R_f=0.3$ was collected and

the solvent was removed under reduced pressure to give white crystalline material (2.377 g, 85% yield). M.p.46-47°C (Lit. m.p.46-47°C).

Reference: Turchin, K.F.; Grokhovskii, S.L.; Zhuze, A.L.; Gottikh, B.P., *Soviet Journal of Bioorganic Chemistry (English Translation)* **1978**, 4, 780-790.

NMR(CDCl₃): 0.97(6H, d, J=6.45Hz), 1.38(3H,t, J=7.2Hz), 1.61-1.73(3H, m), 4.30-4.39(4H, m), 7.43(1H, d, J=1.6Hz), 7.63(1H, d, J=1.6Hz).

IR ν_{\max} [KBr]: 3138, 2970, 1707, 1539, 1510, 1487, 1426, 1389, 1321, 1259, 1204, 1130, 1097 cm⁻¹.

10

(ii) 1-Isopentyl-4-nitro-1H-pyrrole-2carboxylic acid

Ethyl 1-isopentyl-4-nitro-1H-pyrrole-2-carboxylate (2.377 g, 9.329 mmol; see step (i) above) was suspended in ethanol (5 mL), then a solution of NaOH (955 mg dissolved in 20 mL of water) was added. The reaction mixture was reduced by half under reduced pressure at 50°C, and then the cooled solution was extracted with ether. The water layer was cooled to 0°C then conc. HCl was added dropwise with vigorous stirring until pH 2. The white solid material was filtered off, washed with water, dried at 60°C under reduced pressure for 2 h then left at room temperature overnight. The product was obtained as white solid (2.008 g, 95% yield), m.p.154-157°C [Lit.m.p.154-156°C].

20

Reference: Turchin, K.F.; Grokhovskii, S.L.; Zhuze, A.L.; Gottikh, B.P., *Soviet Journal of Bioorganic Chemistry (English Translation)* **1978**, 4, 780-790.

NMR(CDCl₃): 0.98(6H, d, J=6.46Hz), 1.62-1.75(3H, m), 4.38(2H, t, J=7.5Hz), 7.58(1H,d, J=1.98Hz), 7.69(1H, d, J=1.97Hz).

25

IR ν_{\max} [KBr]: 3120, 1677, 1540, 1517, 1480, 1315, 1250 cm⁻¹.

(iii) *N*-[3-(Dimethylamino)propyl]-1-isopentyl-4-nitro-1*H*-pyrrole-2-carboxamide

1-Isopentyl-4-nitro-1*H*-pyrrole-2-carboxylic acid (315 mg, 1.388 mmol; see step (ii) above) was dissolved in thionyl chloride (5 mL) and the reaction mixture heated under reflux for 4 h. Excess thionyl chloride was removed under reduced pressure and the acid chloride so formed was used without further purification. 3-(Dimethylamino)propyl-amine (250 μ L, 2.471 mmol) was dissolved in THF (20 mL, dry) to which NMM (250 μ L, 2.472 mmol) was added. The acid chloride was dissolved in THF (5 mL, dry) then it was added dropwise to the amine solution at room temperature with stirring, which was continued at room temperature overnight. The solvent was removed under reduced pressure and the crude product dissolved in DCM (50 mL) then extracted with aqueous sodium carbonate (540 mg of Na₂CO₃ in 25 mL of water). The organic layer was dried over silica gel using 100:100:1 ethyl acetate / methanol / TFA to give the required product as pale yellow solid (R_f =0.15), (410 mg, 95% yield), m.p.72-73°C.

NMR(CDCl₃): 0.95(6H, d, J =6.5Hz), 1.57-1.76(5H, m), 2.32(6H, s), 2.51(2H,t, J =10.3Hz), 3.47-3.51(2H, qt, J =4.8Hz), 4.40-4.44(2H, q, J =7.5Hz), 6.92(1H,d, J =1.88Hz), 7.56(1H,d, J =1.90Hz), 8.61(1H, s, broad, CONH).

IR ν_{\max} [KBr]: 1656, 1637, 1565, 1534, 1498, 1417, 1333 cm⁻¹.

HREIMS: Found: 310.20031 Calculated for C₁₅H₂₆O₄N₃ 310.20049.

(iv) *N*-[3-(Dimethylamino)propyl]-1-isopentyl-4-{\[(1-isopentyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]amino}-1*H*-pyrrole-2-carboxamide

N-[3-(Dimethylamino)propyl]-1-isopentyl-4-nitro-1*H*-pyrrole-2-carboxamide (151 mg, 0.486 mmol; see step (iii) above) was dissolved in ethanol (20 mL). The reaction mixture was cooled to 0°C then Pd/C-

10%(98 mg) was added to the reaction mixture under N₂. The reaction mixture was then hydrogenated at room temperature and atmospheric pressure for 4 h. The catalyst was removed over Kieselguhr and the solvent removed under reduced pressure at 50°C to give the amine as pale yellow oil that was used in the next step without further purification. 1-Isopentyl-4-nitro-1*H*-pyrrole-2-carboxylic acid (100 mg, 0.441 mmol; see step (ii) above) was dissolved in thionyl chloride (3 mL) and the reaction mixture heated under reflux for 4 h. Excess thionyl chloride was removed under reduced pressure at 50°C and the acid chloride so formed was used without further purification. The amine was dissolved in THF (20 mL, dry) to which NMM (200 µL) was added at room temperature with stirring. The acid chloride was dissolved in THF (15 mL, dry) then added to the amine solution dropwise with stirring at room temperature. The stirring was continued at room temperature overnight. The solvent was removed under reduced pressure at 50°C, then the crude product was dissolved in DCM (50 mL) and extracted with sodium carbonate solution (612 mg in 20 mL of water). The organic layer was collected, dried and the solvent removed under reduced pressure. The crude product was purified by column chromatography using ethyl acetate/methanol/TEA 100:100:1 ratio. Fractions containing the required material were collected and the solvent was then removed under reduced pressure and co-evaporated with *n*-hexane to give the pure product as glassy yellow material $R_f=0.34$, (223 mg, 94% yield), m.p.57-60°C [Transparent].

NMR[CDCl₃]: 0.93(6H, d, J=4.5Hz), 0.95(6H, d, J=4.5Hz), 1.57-1.83(4H, m), 2.35(6H, s), 2.52(2H,t, J=6.1Hz), 3.45-3.49(2H, q, J=5.6Hz), 4.36(2H, t, J=7.4Hz), 4.43(2H, t, J=7.4Hz), 6.51(1H, d, J=1.9Hz), 7.20(1H, d, J=1.9Hz), 7.22(1H, d, J=1.9Hz), 7.64(1H, d, J=1.9Hz), 7.77(2H, s &t).

IR ν_{\max} [KBr]: 2953, 1633, 1592, 1573, 1539, 1525, 1503, 1312 cm⁻¹.

HREIMS: Found: 488.31163 Calculated for C₂₅H₄₀O₄N₆ 488.31110.

(v) *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]-1-isopentyl-4-nitro-1*H*-pyrrole-2-carboxamide

N-[3-(Dimethylamino)propyl]-1-isopentyl-4-{{[(1-isopentyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]amino}-1*H*-pyrrole-2-carboxamide (200 mg, 0.410 mmol; see step (iv) above) was dissolved in ethanol (20 mL) at 0°C. Pd/C-10°C (164 mg) was added to the reaction mixture at 0°C and under N₂. The reaction mixture was hydrogenated at room temperature and atmospheric pressure for 4 h. The catalyst was removed over Kieselguhr and the solvent removed under reduced pressure to give the amine, which was used in the next step without further purification. 1-Isopentyl-4-nitro-1*H*-pyrrole-2-carboxylic acid (110 mg, 0.485 mmol; see step (ii) above) was dissolved in thionyl chloride (3 mL) and heated under reflux for 3 h. Excess thionyl chloride was removed under reduced pressure to give the acid chloride that was used without further purification. The acid chloride was dissolved in THF (20 mL, dry) and added dropwise at room temperature to the amine which was dissolved in THF (20 mL, dry) containing NMM (200 µL). Stirring was continued overnight at room temperature and then the solvent was removed under reduced pressure at 50°C. The residue was dissolved in DCM (50 mL) and extracted with K₂CO₃ (270 mg, in water 20 mL). The organic layer was dried over MgSO₄, filtered and the solvent removed under reduced pressure. The crude product was purified by column chromatography using methanol/ethyl acetate/TEA (100:100:1) to give the product as yellow glassy material (R_f=0.20), 223 mg, 82% yield. M.p. 110-115°C [Transparent].

NMR[CDCl₃]: 0.93(6H, d, J=1.6Hz), 0.95(6H, d, J=5.6Hz), 0.98(6H, d, J=3.7Hz), 1.57-1.78(5H, m), 2.47(6H, s), 2.70(2H, m), 3.49(2H, m), 4.31-4.49(6H, m), 6.57(1H, s), 7.28(1H, s), 7.33(1H, s), 7.41(1H, s), 7.44(1H, s), 7.61(1H, s), 7.64(1H, d, J=1.8Hz), 7.75(1H, s), 8.22(1H, s).

IR ν_{\max} [KBr]: 2953, 1646, 1583, 1534, 1505, 1464, 1422, 1311 cm^{-1} .

HRFABMS: Found: 667.42932 Calculated for $\text{C}_{35}\text{H}_{55}\text{O}_5\text{N}_8$ 667.42954.

(vi) *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]-4-(formylamino)-1-isopentyl-1*H*-pyrrole-2-carboxamide

N-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]-1-isopentyl-4-nitro-1*H*-pyrrole-2-carboxamide (119 mg, 0.178 mmol; see step (v) above) was dissolved in ethanol (20 mL) and Pd/C-10% (104 mg) was added at 0°C under N_2 with stirring. The reaction mixture was hydrogenated at room temperature and atmospheric pressure for 3 h. The catalyst was removed over Kieselguhr, then ethyl formate (15 mL) was added to the ethanolic solution of the amine. The reaction mixture was heated under reflux for 4 days. The solvent was removed under reduced pressure and the crude product was chromatographed over silica gel using ethyl acetate / methanol / TEA (100:100:1). The product was obtained as light brown solid with no distinct m.p., $R_f=0.15$, (84 mg, 71%). Some of this material was purified further by HPLC.

NMR[DMSO- d_6]: 0.87-0.90(18H, m), 1.47-1.53(9H, m), 1.75(2H, m), 2.78(6H, d, $J=4.8\text{Hz}$), 3.06(2H, m), 3.25(2H, m), 4.31(6H, qt, $J=7.2\text{Hz}$), 6.88(1H, d, $J=1.8\text{Hz}$), 6.92(1H, d, $J=1.8\text{Hz}$), 7.02(1H, d, $J=1.8\text{Hz}$), 7.19(1H, d, $J=1.8\text{Hz}$), 7.23(2H, s&d), 8.13(2H, s&t), 9.21(1H, s, broad, TFA), 9.88(1H, s), 9.89(1H, s), 10.05(1H, s).

IR ν_{\max} [KBr]: 2953, 1650, 1580, 1525, 1461, 1402, 1260, 1221 cm^{-1} .

HRFABMS: Found: 665.44877 Calculated for $\text{C}_{36}\text{H}_{57}\text{O}_4\text{N}_8$ 665.45028.

Example 6

N-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopropyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-4-(formylamino)-1-isopropyl-1*H*-pyrrole-2-carboxamide

(i)*N*-[3-(Dimethylamino)propyl]-1-isopropyl-4-nitro-1*H*-pyrrole-2-carboxamide

1-Isopropyl-4-nitro-1*H*-pyrrole-2-carboxylic acid (312 mg, 1.574 mmol; see Example 1, step (iii) above) was dissolved in thionyl chloride (5 mL) and heated until reflux for 4h. The excess thionyl chloride was removed under reduced pressure at 50°C to give the acid chloride as white solid material, which was used without further purification. 3-(Dimethylamino)propylamine (250 µL) was dissolved in THF (20 mL, dry) to which was added NMM (250 µL, dry) at room temperature with stirring. The acid chloride was dissolved in THF (5 mL, dry) and added dropwise to the amine solution at room temperature with stirring. The stirring was continued at room temperature overnight and then the solvent was removed under reduced pressure at 50°C. The crude product was extracted with K₂CO₃ (25 mL, 10%) and DCM (2x50 mL). The organic layer was collected, dried over Na₂SO₄, filtered and the solvent removed under reduced pressure. The crude product obtained as pale yellow oil was purified over silica gel using 49.5/49.5/1% methanol/ethyl acetate/TEA to give the pure product as pale yellow oil, R_F = 0.25, (400 mg, 90% yield).

NMR[CDCl₃]: 8.59(1H, broad); 7.74(1H, d); 6.92(1H, d); 5.65-5.59(1H, qt); 3.51-3.46(2H, q); 2.51(2H, t); 2.32(6H, s); 1.77-1.71(2H, qt); 1.48(6H, d).

IR ν_{max} [NaCl]: 3330, 3136, 2979, 1650, 1535, 1430, 1341, 1279 cm⁻¹.

HREIMS: Found: 282.16908 calculated for C₁₃H₂₂N₄O₃ 282.16919.

(ii) N-[5-({[3-(Dimethylamino)propylamino}carbonyl)-1-isopropyl-1H-pyrrol-3-yl]-1-methyl-4-nitro-1H-pyrrole-2-carboxamide

N-[3-(Dimethylamino)propyl]-1-isopropyl-4-nitro-1H-pyrrole-2-carboxamide (120 mg, 0.425 mmol; see step (i) above) was dissolved in methanol (25 mL) to which Pd/C-10% (52 mg) was added at 0°C under nitrogen. The reaction mixture was hydrogenated at room temperature and atmospheric pressure. The catalyst was removed over Kieselguhr and the solvent was then removed under reduced pressure at 40°C. This gave the amine (as pale yellow oil), which was then dissolved in DCM (25 mL, dry).
1-Methyl-4-nitro-1H-pyrrole-2-carboxylic acid (90 mg, 0.529 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) was suspended in thionyl chloride (4 mL) then heated until reflux for 4 h. The excess thionyl chloride was removed under reduced pressure at 40°C. The acid chloride so obtained was dissolved in DCM (25 mL, dry) then added dropwise with stirring at room temperature to the amine solution. The reaction mixture was left stirring at room temperature overnight. K₂CO₃ solution (232 mg, in water 10 mL) was added and the mixture was extracted with DCM (3x25 mL). The organic layers were combined, dried over MgSO₄, filtered and the solvent removed under reduced pressure. The crude product was purified by flash chromatography using silica gel, methanol/ethyl acetate/TEA [1/3 containing 1%TEA]. The product was obtained as micro-crystalline yellow solid (165 mg, 96%), m.p. 161-163°C.

NMR(CDCl₃): 1.43-1.44(6H, d, J=6.4Hz, 2xMe); 1.72-1.79(2H, quintet, J=6.2Hz, CH₂); 2.33(6H, s, NMe₂); 2.47-2.50(2H, t, J=6.2Hz, CH₂); 3.45-3.49(2H, q, J=5.7Hz, CH₂); 4.04(3H, s, NMe); 5.57-5.62(1H, m); 6.46(1H, d, 1.8J= Hz); 7.22(1H, d, J=1.7Hz); 7.39(1H, d, J=1.7Hz); 7.60(1H, d, J=1.7Hz); 7.73(1H, t, J=5.6Hz, CONH, exch.); 7.77(1H, s, CONH, exch.).

IR ν_{\max} [KBr]: 3281, 3110, 2967, 1640, 1557, 1532, 1503, 1468, 1409, 1310 cm⁻¹.

HRFABMS: Found: 405.22501 Calculated for C₁₉H₂₉N₆O₄ 405.22503.

(iii) *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopropyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-1-isopropyl-4-nitro-1*H*-pyrrole-2-carboxamide

N-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopropyl-1*H*-pyrrol-3-yl]-1-methyl-4-nitro-1*H*-pyrrole-2-carboxamide (160 mg, 0.395 mmol; see step (ii) above) was suspended in methanol (25 mL) to which was added Pd/C-10% (112 mg) at 0°C under nitrogen with stirring. The reaction mixture was hydrogenated at room temperature and atmospheric pressure for 4 h. The catalyst was removed over Kieselguhr and the solvent removed under reduced pressure at 40°C to give the amine. 1-Isopropyl-4-nitro-1*H*-pyrrole-2-carboxylic acid (94 mg, 0.552 mmol; see Example 1, step (iii) above) was suspended in thionyl chloride (4 mL) and heated until reflux for 2 h. Excess thionyl chloride was removed under reduced pressure at 40°C, and the acid chloride so formed was used without further purification. The amine was dissolved in DCM (25 mL, dry) to which TEA (40 µL, dry) was added at room temperature with stirring. The acid chloride was dissolved in DCM (15 mL, dry) after which it was added to the amine solution dropwise with stirring at room temperature. The stirring was continued overnight. Na₂CO₃ solution (280 mg, in water 5 mL) was added to the reaction mixture. The organic layers obtained from the extraction were combined and dried over MgSO₄, filtered and the solvent removed under reduced pressure. The crude product was purified using flash chromatography. Silica gel and methanol/ethyl acetate/TEA [1/3 containing 1% TEA, R_f=0.1] were used to give the product as yellow solid (152 mg, 69%), m.p. 145-148°C.

NMR(DMSO-d₆): 1.32-1.34(6H, d, J=6.4Hz, 2xMe); 1.45-1.46(6H, d, J=6.4Hz, 2xMe); 1.69-1.73(2H, quintet, J=6.2Hz, CH₂); 2.38(6H, s, NMe₂); 2.47-2.50(2H, t, J=6.2Hz, CH₂); 3.19-3.24(2H, m, CH₂); 3.86(3H, s, NMe); 5.42-5.50(1H, m); 6.80(1H, s); 7.05(1H, d, J=1.6Hz); 7.26(1H, d, J=1.6Hz); 7.36(1H, d, J=1.6Hz); 7.50(1H, d, J=1.6Hz); 8.08-8.11(1H, t, J=5.6Hz,

CONH, exch.); 8.33(1H, d, J=1.6Hz); 9.92(1H, s, CONH, exch.); 10.33(1H, s, CONH, exch.).

IR ν_{\max} [KBr]: 2943, 2860, 1638, 1593, 1524, 1456, 1403, 1316, 1251, 1187, 1170 cm^{-1} .

5 HRFABMS: Found: 555.30593 Calculated for $\text{C}_{27}\text{H}_{39}\text{N}_8\text{O}_5$ 555.30434.

(iv) *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopropyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-4-(formyl-amino)-1-isopropyl-1*H*-pyrrole-2-carboxamide

10 *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopropyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-1-isopropyl-4-nitro-1*H*-pyrrole-2-carboxamide (150 mg, 0.291 mmol; see step (iii) above) was dissolved in ethanol (15 mL). The solution was cooled to 0°C under N_2 then Pd/C-10% (120 mg) was added. The reaction mixture was
15 hydrogenated at room temperature and atmospheric pressure for 4 h. The catalyst was removed over Kieselguhr and then ethyl formate (20 mL) was added to the ethanolic solution. The reaction mixture was heated under reflux overnight and then the solvent was removed under reduced pressure at 40°C. The product was purified by HPLC to give the product as white
20 solid material (71 mg, 40% yield) with no distinct melting point.

NMR[DMSO- d_6]: 10.0564(1H, s); 9.96(1H,s); 9.90(1H, s); 9.24(1H,s, broad, TFA); 8.16-8.14(2H, m); 7.34(2H, s); 7.22(1H, s); 7.06(1H, s); 6.89(1H, s); 6.68(1H, s); 5.49-5.42(1H, m); 3.84(3H, s); 3.25(2H, m); 3.06(2H, m); 2.79(6H, d, J=4.76Hz); 1.84(2H, m); 1.36-1.33(12H, m).

25 IR ν_{\max} [KBr]: 3425, 3285, 2972, 1676, 1639, 1579, 1533, 1465, 1439, 1404, 1257, 1200, 1180, cm^{-1} .

HRFABMS: Found 554.33484 calculated for $\text{C}_{28}\text{H}_{42}\text{N}_8\text{O}_4$ 554.33290

Example 7

N-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-2-({[4-(formylamino)-1-methyl-1*H*-pyrrol-2-yl]carbonyl}amino)-5-isopropyl-1,3-thiazole-4-carboxamide

5

(i) *N*-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-5-isopropyl-2-{{[1-methyl-4-nitro-1*H*-pyrrol-2-yl]carbonyl}amino}-1,3-thiazole-4-carboxamide

N-[3-(Dimethylamino)propyl]-1-methyl-4-nitro-1*H*-pyrrole-2-carboxamide
10 (270 mg, 1.062 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) was dissolved in methanol (25 mL), to which Pd/C-10% (236 mg) was added at 0°C under N₂ with stirring. The reaction mixture was hydrogenated at room temperature and atmospheric pressure for 4 h after which the catalyst was removed over Kieselguhr and the solvent was then removed under reduced
15 pressure to give the amine. 5-Isopropyl-2-{{[1-methyl-4-nitro-1*H*-pyrrol-2-yl]carbonyl}amino}-1,3-thiazole-4-carboxylic acid (359 mg, 1.062 mmol; see Example 3, step (iii) above) was dissolved in DMF (4 mL, dry) to which NMM (0.5 mL, dry) was added followed by HBTU (721 mg, 1.901 mmol). The amine was dissolved in DMF (2 mL, dry) and then added to the
20 reaction mixture at room temperature with stirring. The stirring was continued at room temperature overnight. NMM was removed under reduced pressure. The reaction mixture was diluted with ethyl acetate then NaHCO₃ saturated was added with stirring. The organic layer was separated, dried over MgSO₄ and the solvent was removed under reduced
25 pressure. The crude product obtained was chromatographed over silica gel using 1:1:0.2 methanol/ethyl acetate/TEA. The product was obtained as yellow glassy material (415 mg, 72% yield) with no distinct melting point. NMR[DMSO-*d*₆]: 9.60(1H,s); 8.23(1H, s); 8.09-8.036(1H, t); 7.83(1H, s); 7.25(1H, d); 6.91(1H, d); 4.22-4.16(1H, qt); 4.00(3H, s); 3.82(3H, s); 3.25-

3.18(2H, q); 2.38-2.34(2H, t); 2.23(6H, s); 1.68-1.61(2H, qt); 1.30-1.28(6H, d).

IR ν_{\max} [KBr]: 1644, 1551, 1500, 1465, 1402, 1309, 1283, 1194, 1100 cm^{-1} .

HRFABMS: Found: 545.22962 calculated for $\text{C}_{24}\text{H}_{33}\text{N}_8\text{O}_5\text{S}$ 545.22946.

5

(ii) *N*-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-2-({[4-(formylamino)-1-methyl-1*H*-pyrrol-2-yl]carbonyl}-amino)-5-isopropyl-1,3-thiazole-4-carboxamide

N-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-5-isopropyl-2-{{[(1-methyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]amino}-1,3-thiazole-4-carboxamide (137 mg, 0.252 mmol; see step (i) above) was dissolved in ethanol (15 mL), to which Pd/C-10% (121 mg) was added under N_2 at 0°C. The reaction mixture was hydrogenated at room temperature and atmospheric pressure for 2.5 h. The catalyst was removed over Kieselguhr and then ethyl formate (25 mL) was added to the ethanolic solution. The reaction mixture was heated under reflux overnight, following which the solvent was removed under reduced pressure and the product was then purified by HPLC. The product was obtained after freeze-drying as white solid material (41 mg, 25% yield) [TFA salt] with no distinct melting point.

15
20

NMR[DMSO- d_6]: 12.0971(1H, s); 10.1404(1H, s); 9.5957(1H, s); 9.2310(1H, broad, TFA); 8.3533-8.3249(1H, d and 1H, t); 7.3982(1H, s); 7.2894(1H, d); 7.2046(1H, d); 7.0146(1H, d); 4.1717-4.1200(1H, m); 3.8841(3H, s); 3.8274(3H, s); 3.2577(2H, m); 3.0646(2H, m); 2.7987(6H, d, NMe₂); 1.8612-1.8231(2H, m); 1.3135(6H, d).

25

IR ν_{\max} [KBr]: 3428, 1662, 1545, 1467, 1401, 1284, 1201, 1134 cm^{-1} .

HRFABMS: Found: 543.24988 calculated for $\text{C}_{25}\text{H}_{35}\text{N}_8\text{O}_4\text{S}$ 543.25020.

Example 8

4-({[4-(Formylamino)-1-methyl-1*H*-pyrrol-2-yl]carbonyl}amino)-1-iso-propyl-*N*-[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1*H*-pyrrole-2-carboxamide

5

(i) 1-Isopropyl-*N*-[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]-4-nitro-1*H*-pyrrole-2-carboxamide

1-Methyl-*N*-[3-(4-morpholinyl)propyl]-4-nitro-1*H*-pyrrole-2-carboxamide (214 mg, 0.805 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) was dissolved in methanol (25 mL). The solution was cooled to 0°C under N₂ then Pd/C-10% (192 mg) was added. The reaction mixture was hydrogenated at room temperature and atmospheric pressure for 3 h. The catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure at 50°C. The amine so formed was dissolved in DMF (2 mL, dry), to which 1-isopropyl-4-nitro-1*H*-pyrrole-2-carboxylic acid (160 mg, 0.805 mmol; see Example 1, step (iii) above) was added followed HBTU (610 mg, 1.608 mmol) and NMM (100 µL, dry). Additional amount of DMF (2 mL, dry) was added with stirring at room temperature and was continued for 72 h. The reaction mixture was diluted with ethyl acetate and NaHCO₃ solution with stirring. The organic layer was separated, dried MgSO₄, filtered and the solvent removed under reduced pressure. The crude product was chromatographed over flash silica gel using 1/3 methanol/ethyl acetate. Fractions containing the product [R_F=0.30] were collected and the solvents were removed under reduced pressure at 50°C. The product was obtained as glassy yellow material [hygroscopic] (260 mg, 73% yield) with no distinct melting point.

NMR[DMSO-*d*₆]: 10.28(1H, s); 8.33(1H, d); 8.08-8.05(1H, t); 7.48(1H, d); 7.20(1H, d); 6.85(1H, d); 5.89-5.42(1H, qt); 3.80(3H, s); 3.58-3.56(4H, t); 3.22-3.17(2H, q); 2.32-2.28(6H, m); 1.67-1.62(2H, qt); 1.45-1.43(6H, d).

IR ν_{\max} [KBr]: 1661, 1582, 1540, 1441, 1389, 2326, 1284, 1242, 1116, 850, 755 cm^{-1} .

HREIMS: found 446.22568 calculated for $\text{C}_{21}\text{H}_{30}\text{N}_6\text{O}_5$ 446.22777.

- 5 (ii) 1-Isopropyl-*N*-[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}-carbonyl)-1*H*-pyrrol-3-yl]-4-{{[(1-methyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]-amino}-1*H*-pyrrole-2-carboxamide

1-Isopropyl-*N*-[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]-4-nitro-1*H*-pyrrole-2-carboxamide (250 mg, 0.560 mmol;
10 see step (i) above) was dissolved at 0°C under N_2 . Pd/C-10% (220 mg) was added and the reaction mixture was hydrogenated for 3.5 h. The catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure. The amine so formed was dissolved in DMF (2 mL, dry), to which was NMM (100 μL , dry) was added followed by HBTU (425 mg,
15 1.069 mmol) and 1-methyl-4-nitro-1*H*-pyrrole-2-carboxylic acid (95 mg, 0.560 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) with stirring at room temperature. The stirring was continued overnight. The reaction mixture was diluted with ethyl acetate and a saturated solution of sodium hydrogen carbonate. The organic layer was separated, dried over MgSO_4 and filtered.
20 The solvent was removed under reduced pressure and the crude product was chromatographed using flash silica gel and 1:3:0.1 methanol/ethyl acetate/TEA. Fractions containing the pure material were collected [$R_F=0.3$] solvents were removed under reduced pressure at 40°C to give semi-solid glassy yellow material (170 mg, 53%) with no distinct melting
25 point.

NMR[DMSO-d_6]: 10.31(1H,s); 9.98(1H,s); 8.20(1H,s); 8.04-8.01(1H,t); 7.59(1H,s); 7.44(1H,s); 7.18(1H,s); 6.92(1H,s); 6.86(1H,s); 5.49-5.42(1H,qt); 3.97(3H,s); 3.80(3H,s); 3.58-3.57(4H,t); 3.22-3.17(2H,q); 2.33-2.29(2H,d); 1.68-1.61(2H,qt); 1.38-1.36(6H,d); 1.05-1.01(2H,m).

IR ν_{\max} [KBr]: 3425, 2927, 1662, 1653, 1582, 1543, 1502, 1410, 1307, 1251, 1203, 1115, 848 cm^{-1} .

HRFABMS: found 569.28147 calculated for $\text{C}_{27}\text{H}_{37}\text{N}_8\text{O}_6$ 569.28361.

- 5 (iii) 4-({[4-(Formylamino)-1-methyl-1*H*-pyrrol-2-yl]carbonyl}amino)-1-isopropyl-*N*-[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1*H*-pyrrole-2-carboxamide

1-Isopropyl-*N*-[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]-4-{{[(1-methyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]amino}-1*H*-pyrrole-2-carboxamide (160 mg, 0.282 mmol; see step (ii) above) was dissolved in ethanol (25 mL) to which Pd/C-10% (150 mg) was added at 0°C under nitrogen with stirring. The reaction mixture was hydrogenated for 2 h at room temperature and atmospheric pressure. The catalyst was removed over Kieselguhr. Ethyl formate (25 mL) was added to the ethanolic solution, then the reaction mixture was heated under reflux overnight. Ethanol and excess ethyl formate were removed under reduced pressure and the crude product was purified by HPLC to give the required material as white solid (40.1 mg, 21% yield) with no distinct melting point.

15 NMR[DMSO- d_6]: 10.05(1H,s); 9.93(1H,s); 9.92(1H,s); 9.58(1H, broad, TFA); 8.17(1H,t); 8.13(1H,s); 7.39(1H,s); 7.19(1H,s); 7.17(1H,s); 6.97(1H,s); 6.95(1H,s); 6.92(1H,s); 3.84(3H,s); 3.81(3H,s); 3.56-3.60(4H,t); 3.49-3.43(4H,t); 3.26-3.24(2H,q); 3.12-3.06(4H,m); 1.87(2H,qt); 1.37-1.36(6H,d).

IR ν_{\max} [KBr]: 3423, 1672, 1644, 1581, 1536, 1462, 1440, 1407, 1254, 1200, 1134 cm^{-1} .

HRFABMS: found: 567.30312 calculated for $\text{C}_{28}\text{H}_{39}\text{N}_8\text{O}_5$ 567.30434.

Example 9

4-(Formylamino)-N-[1-isopropyl-5-({[1-methyl-5-({[3-(1-pyrrolidinyl)-propyl]amino}carbonyl)-1H-pyrrol-3-yl]amino}carbonyl)-1H-pyrrol-3-yl]-1-methyl-1H-pyrrole-2-carboxamide

5

(i) 1-Methyl-4-nitro-N-[3-(1-pyrrolidinyl)propyl]-1H-pyrrole-2-carboxamide

3-(1-Pyrrolidino)propylamine (236 mg, 1.842 mmol (Lancaster Synthesis Ltd.)) was dissolved in THF (25 mL, dry) to which a solution of 2,2,2-trichloro-1-(1-methyl-4-nitro-1H-pyrrol-2-yl)ethanone (508 mg, 1.842 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) was added in THF (2 mL, dry) at room temperature with stirring. The reaction mixture was left stirring at room temperature overnight. The solvent was removed under reduced pressure and the crude product was purified by column chromatography using 49.5:49.5:1 ethyl acetate/ methanol/ TEA. The product was obtained as glassy yellow solid, $R_f = 0.1$ (503 mg, 98% yield), m.p. 113-115°C.

$^1\text{H-NMR}$ (DMSO- d_6) δ 1.70-1.79(6H, m), 2.48-2.51(6H, m), 3.28-3.51(2H, q, $J=6.85\text{Hz}$), 3.89(3H, s), 7.45(1H, d, $J=1.6\text{Hz}$), 8.18(1H, d, $J=1.6\text{Hz}$), 8.51(1H, t, $J=5.37\text{Hz}$).

IR ν_{max} [KBr]: 3131, 2966, 2794, 1654, 1545, 1525, 1492, 1305 cm^{-1} .

HRFABMS: Found: 281.16249 Calculated for $\text{C}_{13}\text{H}_{21}\text{N}_4\text{O}_3$ 281.16137.

(ii) 1-Isopropyl-N-[1-methyl-5-({[3-(1-pyrrolidinyl)propyl]amino}-carbonyl)-1H-pyrrol-3-yl]-4-nitro-1H-pyrrole-2-carboxamide

25

1-Methyl-4-nitro-N-[3-(1-pyrrolidinyl)propyl]-1H-pyrrole-2-carboxamide (227 mg, 0.811 mmol; see step (i) above) was dissolved in methanol (25 mL) to which Pd/C-10% (160 mg) was added at 0°C under nitrogen with stirring. The reaction mixture was hydrogenated for 2 h at room

temperature and atmospheric pressure. The catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure at 40°C to give the amine. 1-Isopropyl-4-nitro-1*H*-pyrrole-2-carboxylic acid (161 mg, 0.811 mmol; see Example 1, step (iii)) was dissolved in DMF (1 mL, dry) to which HBTU (615 mg, 1.622 mmol) and NMM (100 µL, dry) were added. The amine was dissolved in DMF (2 mL, dry) and added to the reaction mixture at room temperature with stirring. The stirring was continued at room temperature overnight. The reaction mixture was diluted with ethyl acetate and a saturated solution of sodium hydrogen carbonate. The organic layer was separated, dried over magnesium sulphate, filtered and the solvent removed under reduced. The product was purified by column chromatography using 1/2/0.1 methanol/ethyl acetate/TEA. The product was obtained as yellow material (300 mg, 86% yield) with no distinct melting point. Some of this material was further purified by HPLC.

NMR[DMSO- d_6]: 10.31(1H, s); 9.48(1H, broad, TFA); 8.34(1H, d); 8.21-8.18(1H, t); 7.48(1H, d); 7.28(1H, d); 6.92(1H, d); 5.48-5.42(1H, qt); 3.82(3H, s); 3.56-3.55(2H, m); 3.27-3.23(2H, q); 3.17-3.12(2H, m); 2.99-2.97(2H, m); 2.01(2H, m); 1.86-1.83(4H, m); 1.45-1.44(6H, d).

IR ν_{\max} [KBr]: 3433, 1672, 1650, 1535, 1462, 1426, 1371, 1321, 1283, 1237, 1200, 1178, 1131 cm^{-1} .

HRFABMS: found: 430.23179 calculated for $\text{C}_{21}\text{H}_{30}\text{N}_6\text{O}_4$ 430.23285.

(iii) *N*-[1-Isopropyl-5-({[1-methyl-5-({[3-(1-pyrrolidinyl)propyl]amino}-carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1-methyl-4-nitro-1*H*-pyrrole-2-carboxamide

1-Isopropyl-*N*-[1-methyl-5-({[3-(1-pyrrolidinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]-4-nitro-1*H*-pyrrole-2-carboxamide (300 mg, 0.697 mmol; see step (ii) above) was dissolved in methanol (25 mL) to which Pd/C-10% (235 mg) was added at 0°C under nitrogen with stirring. The reaction

mixture was hydrogenated for 3 h at room temperature and atmospheric pressure. The catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure at 40°C to give the amine. 1-Methyl-4-nitro-1*H*-pyrrole-2-carboxylic acid (119 mg, 0.699 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) was dissolved in DMF (2 mL, dry) to which HBTU (529 mg, 1.394 mmol) and NMM (100 µL, dry) were added. The amine was dissolved in DMF (2 mL, dry) and added to the reaction mixture at room temperature with stirring. The stirring was continued at room temperature overnight. The reaction mixture was diluted with ethyl acetate and a saturated solution of sodium hydrogen carbonate. The organic layer was separated, dried over magnesium sulphate, filtered and the solvent removed under reduced pressure. The product was purified by column chromatography using 1/2/0.1 methanol/ethyl acetate/TEA. The product was obtained as yellow material (239 mg, 62% yield) with no distinct melting point. Some of this material was further purified by HPLC.

NMR[DMSO- d_6]: 10.28(1H,s); 9.98(1H,s); 9.46(1H, broad, TFA); 8.19(1H,s); 8.14(1H,t); 7.58(1H,s); 7.43(1H,s); 7.18(1H,s); 6.94(2H,s); 5.46-5.43(1H,m); 3.97(3H,s); 3.82(3H,s); 3.55(2H,m); 3.27(2H,m); 3.14(2H,m); 3.00(2H,m); 2.02(2H,m); 1.86(4H,m); 1.38-1.37(6H,d).

IR ν_{\max} [KBr]: 3419, 3136, 1673, 1645, 1586, 1531, 1463, 1409, 1311, 1247, 1199, 1130 cm^{-1} .

HRFABMS: found: 553.29109 calculated for $\text{C}_{27}\text{H}_{37}\text{N}_8\text{O}_5$ 553.28869.

(iv) 4-(Formylamino)-*N*-[1-isopropyl-5-({[1-methyl-5-({[3-(1-pyrrolidinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1-methyl-1*H*-pyrrole-2-carboxamide

N-[1-Isopropyl-5-({[1-methyl-5-({[3-(1-pyrrolidinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1-methyl-4-nitro-1*H*-pyrrole-2-carboxamide (224 mg, 0.406 mmol; see step (iii) above)

was dissolved in ethanol (25 mL) to which Pd/C-10% (224 mg) was added at 0°C under nitrogen with stirring. The reaction mixture was hydrogenated for 2.5 h at room temperature and atmospheric pressure. The catalyst was removed over Kieselguhr. Ethyl formate (15 mL) was added to the ethanolic solution, then the reaction mixture was heated under reflux overnight. Ethanol and excess ethyl formate were removed under reduced pressure and the crude product was purified by HPLC to give the required material as white solid (42.7 mg, 16% yield) with no distinct melting point. NMR[DMSO-d₆]: 10.07(1H,s); 9.94(1H,s); 9.92(1H,s); 9.42(1H, broad, TFA); 8.17-8.13(1H,2H,m); 7.39(1H,d); 7.19(1H,d); 7.17(1H,d); 6.97(1H,d); 6.95(1H,d); 6.92(1H,d); 3.84(3H,s); 3.61(3H,s); 3.55(2H,m); 3.26-3.24(2H,m); 3.17-3.12(2H,m); 3.00(2H,m); 2.01(2H,m); 1.86(4H,m); 1.37-1.36(6H,d). IR ν_{\max} [KBr]: 3429, 1675, 1645, 1582, 1534, 1462, 1440, 1407, 1252, 1200, 1132 cm⁻¹. HRFABMS: found: 551.31199 calculated for C₂₈H₃₉N₈O₄ 551.30943.

Example 10

N-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1H-pyrrol-3-yl]-4-(formylamino)-1-methyl-1H-pyrrole-2-carboxamide

(i) N-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]-1-isopentyl-4-nitro-1H-pyrrole-2-carboxamide
N-[3-(Dimethylamino)propyl]-1-methyl-4-nitro-1H-pyrrole-2-carboxamide (253 mg, 0.995 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) was dissolved in methanol (25 mL) to which Pd/C-10% (162 mg) was added at 0°C under nitrogen. The reaction mixture was hydrogenated for 4 h. The catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure at 50°C to give the amine. 1-Isopentyl-4-nitro-1H-

pyrrole-2-carboxylic acid (228 mg, 1.005 mmol; prepared according to the procedure described in *Soviet Journal of Bioorganic Chemistry (English Translation)* 4, 780-790 (1978)) was dissolved in thionyl chloride (4 mL) and the reaction mixture was heated under reflux for 4 h. Excess thionyl chloride was removed under reduced pressure and the acid chloride so formed was dissolved in DCM (10 mL, dry). The amine was dissolved in DCM (10 mL, dry) to which NMM (200 μ L, dry) was added at room temperature with stirring. The acid chloride solution was added dropwise to the amine solution at room temperature with stirring. The stirring was continued overnight at room temperature. The reaction mixture was diluted with DCM (25 mL) and saturated solution of sodium hydrogen carbonate (25 mL). After the extraction, the organic layer was collected, dried over magnesium sulphate, filtered and the solvent removed under reduced pressure. The product was purified by column chromatography using 1/3/0.030 methanol/ethyl acetate/TEA. Fractions containing the product ($R_F=0.20$) were collected and the solvents were removed under reduced pressure at 50°C to give the product as yellow powder (347 mg, 81% yield), m.p.148-151°C.

NMR[DMSO- d_6]: 10.23(1H,s); 8.23(1H,d); 8.14-8.11(1H,t); 7.55(1H,d); 7.19(1H,d); 6.82(1H,d); 4.45-4.41(2H,t); 3.81(3H,s); 3.22-3.17(2H,q); 2.35-2.31(2H,t); 2.20(6H,s); 1.67-1.58(4H,m); 1.56-1.46(1H,m); 0.89-0.87(6H,d).

IR ν_{max} [KBr]: 1665, 1642, 1599, 1528, 1499, 1425, 1312 cm^{-1} .

HRFABMS: found 433.25630 calculated for $C_{21}H_{33}N_6O_4$ 433.25633.

(ii) N-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1H-pyrrol-3-yl]-1-methyl-4-nitro-1H-pyrrole-2-carboxamide

N-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]-1-isopentyl-4-nitro-1H-pyrrole-2-carboxamide (335 mg, 0.775 mmol; see step (i) above) was dissolved in methanol (25 mL) to which Pd/C-10% (213 mg) was added at 0°C under nitrogen. The reaction mixture was hydrogenated for 5 h. The catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure at 50°C to give the amine. 1-Methyl-4-nitro-1H-pyrrole-2-carboxylic acid (210 mg, 1.234 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) was dissolved in thionyl chloride (3 mL) and the reaction mixture was heated under reflux for 5 h. Excess thionyl chloride was removed under reduced pressure and the acid chloride so formed was dissolved in DCM (10 mL, dry). The amine was dissolved in DCM (10 mL, dry) to which NMM (200 µL, dry) was added at room temperature with stirring. The acid chloride solution was added dropwise to the amine solution at room temperature with stirring. The stirring was continued overnight at room temperature. The reaction mixture was diluted with DCM (25 mL) and saturated solution of sodium hydrogen carbonate (25 mL). After the extraction, the organic layer was collected, dried over magnesium sulphate, filtered and the solvent removed under reduced pressure. The product was purified by column chromatography using 1/3/0.030 methanol/ethyl acetate/TEA. Fractions containing the product ($R_F=0.20$) were collected and the solvents were removed under reduced pressure at 50°C to give the product as yellow solid material (335 mg, 78% yield), m.p. 110-115°C [transparent].

NMR[DMSO- d_6]: 10.28(1H,s); 9.92(1H,s); 8.18(1H,d); 8.10-8.07(1H,t); 7.59(1H,d); 7.32(1H,d); 7.18(1H,d); 6.98(1H,d); 6.91(1H,d); 6.83(1H,d); 4.35-4.32(2H,t); 3.97(3H,s); 3.80(3H,s); 3.22-3.17(2H,q); 2.36-2.32(2H,t);

2.21(6H,s); 1.68-1.62(2H,q); 1.61-1.45(3H,m); 1.05-0.99(2H,m); 0.90-0.88(6H,d).

IR ν_{\max} [KBr]: 3401, 3289, 1643, 1582, 1533, 1594, 1463, 1400, 1309, 1253 cm^{-1} .

5 HRFABMS: found 555.30169 calculated for $\text{C}_{27}\text{H}_{39}\text{N}_8\text{O}_5$ 555.30434.

(iii) *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]-4-(formylamino)-1-methyl-1*H*-pyrrole-2-carboxamide

10 *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]-1-methyl-4-nitro-1*H*-pyrrole-2-carboxamide (89 mg, 0.160 mmol; see step (ii) above) was dissolved in ethanol (25 mL) to which Pd/C-10% (95 mg) was added at 0°C under nitrogen with stirring. The reaction mixture was hydrogenated for 2 h
15 at room temperature and atmospheric pressure. The catalyst was removed over Kieselguhr. Ethyl formate (25 mL) was added to the ethanolic solution, then the reaction mixture was heated under reflux overnight. Ethanol and excess ethyl formate were removed under reduced pressure and the crude product was purified by HPLC to give the required material as
20 pale yellow solid (37.4 mg, 35% yield) with no distinct melting point.

NMR[DMSO- d_6]: 10.04(1H,s); 9.90(1H,s); 9.88(1H,s); 9.63(1H, broad, TFA); 8.15-8.13(2H,s & t); 7.26(1H,s); 7.18(1H,s); 7.16(1H,s); 7.01(1H,s); 6.92(1H,s); 6.92(1H,s); 4.31-4.30(2H,t); 3.84(3H,s); 3.82(3H,s); 3.25-3.24(2H,m); 3.07(2H,m); 2.79(6H,d); 1.84-1.82(2H,m); 1.57-1.48(3H,m);
25 0.90(6H,d).

IR ν_{\max} [KBr]: 3428, 3315, 1676, 1644, 1582, 1538, 1464, 1438, 1403, 1262, 1201, 1132 cm^{-1} .

HRFABMS: found 553.32520 calculated for $\text{C}_{28}\text{H}_{41}\text{N}_8\text{O}_4$ 553.32508.

Example 11

2-(Acetylamino)-N-[5-({[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]-5-isopropyl-1,3-thiazole-4-carboxamide

5

(i) Methyl 2-(acetylamino)-5-isopropyl-1,3-thiazole-4-carboxylate

Methyl 2-amino-5-isopropyl-1,3-thiazole-4-carboxylate (500 mg, 2.497 mmol; see Example 3, step (i) above) was dissolved in DCM (10 mL, dry) to which NMM (300 μ L, dry) was added at room temperature with stirring. Acetyl chloride (300 μ L, 330 mg, 4.204 mmol) was added dropwise to the reaction mixture with stirring. The stirring was continued at room temperature overnight. Brine (25 mL) was added to the reaction mixture and the product was then extracted with DCM (2x50 mL). The organic layers were combined and dried over MgSO_4 , filtered and the solvent removed under reduced pressure to give the crude product which was purified by column chromatography over silica gel using 1/1 ethyl acetate / n-hexane [R_F =0.22]. The product was obtained as white fine needles (432 mg, 72% yield), m.p. 170-172°C.

NMR(DMSO- d_6): 12.31(1H, s); 4.01-3.94(1H, quintet); 3.78(3H, s); 2.11(3H, s); 1.27(6H, d).

IR ν_{max} [KBr]: 3277, 2971, 1752, 1694, 1543, 1202 cm^{-1} .

HRFABMS: Found 243.08074 calculated for $\text{C}_{10}\text{H}_{15}\text{N}_2\text{O}_3\text{S}$ 243.08034.

(ii) 2-(Acetylamino)-5-isopropyl-1,3-thiazole-4-carboxylic acid

Methyl 2-(acetylamino)-5-isopropyl-1,3-thiazole-4-carboxylate (242 mg, 1.00 mmol; see step (i) above) was dissolved in ethanol (10 mL) to which ethanolic sodium hydroxide [0.5 M, 10 mL] was added. The reaction mixture was heated under reflux for 30 min following which it was cooled to 0°C and then dilute HCl was added with stirring until pH 2. Water

(50 mL) was added and the white precipitate was filtered off and dried under reduced pressure at 45°C overnight. The product was obtained as micro-crystalline white solid (166 mg, 73% yield), m.p. >230°C.

NMR(DMSO- d_6): 12.65(1H, br); 12.23(1H, s); 4.04-3.97(1H, quintet);
5 2.11(3H, s); 1.26(6H, d).

IR ν_{\max} [KBr]: 3076, 2971, 2868, 1699, 1671, 1568, 1543, 1301, 1221 cm^{-1} .

HRFABMS: Found 229.06408 calculated for $\text{C}_9\text{H}_{13}\text{N}_2\text{O}_3\text{S}$ 229.06469.

(iii) 2-(Acetylamino)-N-[5-({[5-({[3-(dimethylamino)propyl]amino}-
10 carbonyl)-1-methyl-1H-pyrrol-3-yl]amino}carbonyl)-1-methyl-1H-pyrrol-
3-yl]-5-isopropyl-1,3-thiazole-4-carboxamide

N-[5-({[3-Dimethylamino]propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]-1-methyl-4-nitro-1H-pyrrole-2-carboxamide (105 mg, 0.279 mmol; see *Tetrahedron* 56, 5225-5239, (2000)) was dissolved in methanol (25 mL).

15 The solution was cooled to 0°C under N_2 then Pd/C-10% (82 mg) was added. The reaction mixture was hydrogenated at room temperature and atmospheric pressure for 3 h. The catalyst was removed over Kieselguhr and then the solvent was removed under reduced pressure at 50°C. The amine thus formed was dissolved in DMF (2 mL, dry), to which 2-
20 (acetylamino)-5-isopropyl-1,3-thiazole-4-carboxylic acid (96 mg, 0.418 mmol; see step (ii) above) was added, followed by HBTU (211 mg, 0.556 mmol) and NMM (300 μL , dry). The reaction mixture was left stirring at room temperature overnight after which it was diluted with ethyl acetate and NaHCO_3 solution with stirring. The organic layer was
25 separated, dried MgSO_4 , filtered and the solvent removed under reduced pressure. The product was purified by HPLC to give the pure material as white solid (71.3 mg, 38% yield) with no distinct melting point.

NMR[DMSO- d_6]: 12.31(1H, s); 8.87(1H, s); 9.57(1H, s); 9.24(1H, broad); 8.14(1H, t); 7.26(1H, d); 7.17(1H, d); 7.12(1H, d); 6.94(1H, d); 4.19-

4.12(1H, m); 3.85(3H, s); 3.81(3H, s); 3.25(2H, m); 3.07(2H, m); 2.79(6H, d); 2.16(3H, s); 1.83(2H, qt); 1.29(6H, d).

IR ν_{\max} [KBr]: 1676, 1644, 1550, 1465, 1435, 1404, 1202, 1136 cm^{-1} .

HRFABMS: found 557.26442 calculated for $\text{C}_{26}\text{H}_{37}\text{N}_8\text{O}_4\text{S}$ 557.26585.

5

Example 12

2-(Acetylamino)-N-[5-({[4-({[3-(dimethylamino)propyl]amino}carbonyl)-5-isopropyl-1,3-thiazol-2-yl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-5-isopropyl-1,3-thiazole-4-carboxamide

- 10 *N*-[3-(Dimethylamino)propyl]-5-isopropyl-2-[(1-methyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]amino}-1,3-thiazole-4-carboxamide (100 mg, 0.236 mmol; see Example 3, step (iv) above) was dissolved in methanol (25 mL). The solution was cooled to 0°C under N_2 then Pd/C-10% (82 mg) was added. The reaction mixture was hydrogenated at room temperature
- 15 and atmospheric pressure for 3 h. The catalyst was removed over Kieselguhr and the solvent was then removed under reduced pressure at 50°C. The amine so formed was dissolved in DMF (2 mL, dry), to which 2-(acetylamino)-5-isopropyl-1,3-thiazole-4-carboxylic acid (64 mg, 0.356 mmol; see Example 11, step (ii) above) was added, followed by
- 20 HBTU (135 mg, 0.356 mmol) and NMM (200 μL , dry). The reaction mixture was left stirring at room temperature overnight, after which it was diluted with ethyl acetate and NaHCO_3 solution with stirring. The organic layer was separated, dried MgSO_4 , filtered and the solvent removed under reduced pressure. The product was purified by HPLC to give the pure
- 25 material as white solid (78.8 mg, 47% yield) with no distinct melting point. NMR[$\text{DMSO}-d_6$]: 12.11(1H, s); 12.01(1H, s); 9.64(1H, s); 9.27(1H, broad); 7.96(1H, t); 7.51(1H, d); 7.42(1H, d); 4.22-4.11(2H, m); 3.90(3H, s); 3.34(2H, m); 3.08(2H, m); 2.79(6H, d); 2.16(3H, s); 1.91-1.85(2H, qt); 1.30-1.27(12H, 2xd).

IR ν_{\max} [KBr]: 1666, 1549, 1508, 1466, 1398, 1286, 1201, 1134 cm^{-1} .

HRFABMS: found 603.25162 calculated for $\text{C}_{27}\text{H}_{39}\text{N}_8\text{O}_4\text{S}_2$ 603.25357.

Example 13

5 2-(Acetylamino)-N-(5-{[(3-{[3-(dimethylamino)propyl]amino}-3-oxo-propyl)amino]carbonyl}-1-methyl-1*H*-pyrrol-3-yl)-5-isopropyl-1,3-thiazole-4-carboxamide

Samples of Boc-BALa-PAM resin (100 mg of 100-200 mesh (Novabiochem)) were placed in 10 separate vials to each of which was added DCM (0.5 mL, dry). TFA (80%) in DCM (3 mL, dry) was then added to each vial. The reactions were carried out under nitrogen and the reaction mixture agitated for 15 min. Following this, the solvents were drained and the resin washed with DMF (2 x 2 mL, dry) and they were then agitated for 5 min and then drained. 4-[(*tert*-Butoxycarbonyl)amino]-1-methyl-1*H*-pyrrole-2-carboxylic acid (180 mg, 0.750 mmol; see *J. Med. Chem.* 46(17), 3492-3497 (1981)) was dissolved in DMF (15 mL, dry) and then (1.5 mL) of this solution was added to each vial. HBTU (284 mg, 0.748 mmol) was dissolved in DMF (15 mL, dry) then (1.5 mL) of this solution was added to each vial followed by diisopropylethylamine (0.5 mL, dry). The agitation was continued for 30 min after which the solvents and reagents were drained. TFA (80%) in DCM (3 mL, dry) were added to each vial. The reaction mixture was agitated for a further 15 min. The solvents were drained and the resin was then washed with DMF (2 x 2 mL, dry) and agitated for 5 min and then drained. 2-(Acetylamino)-5-isopropyl-1,3-thiazole-4-carboxylic acid (171 mg, 0.750 mmol; see Example 11, step (ii) above) was dissolved in DMF (15 mL, dry) then (1.5 mL) of this solution was added to each vial. HBTU (284 mg, 0.748 mmol) was dissolved in DMF (15 mL, dry) then 1.5 mL of this solution, followed by diisopropylethylamine (0.5 mL, dry), was added to each vial. Agitation was continued for 30 min after which the solvents and reagents were drained.

The resin in each vial was washed successively with DMF (3 mL, dry) for 5 min with agitation, drained and then followed by DCM (3 mL, dry), drained again and finally washed with THF (3 mL, dry) and drained again. To each vial THF (3 mL, dry) was added followed by 3-(dimethylamino)-propylamine (0.5 mL, dry). Agitation was continued overnight at 55°C, then the solvent was filtered. The combined filtrate was collected and the solvent and reagent were removed under reduced pressure to give the crude product as brown oil. This material was further purified by HPLC and the fractions containing the required material were freeze-dried. The pure product was obtained as white solid (92 mg, 20% yield) with no distinct melting point.

NMR[DMSO-d₆]: 12.07(1H, s); 9.51(1H, s); 9.24(1H, broad); 8.03(2H, t); 7.21(1H, d); 6.91(1H, d); 4.17-4.11(1H, m); 3.81(3H, s); 3.41-3.38(2H, m); 3.13-3.09(2H, m); 3.01(2H, m); 2.74(6H, t); 2.35(2H, t); 2.15(3H, s); 1.77-1.71(2H, qt); 1.28(6H, d).

IR ν_{\max} [KBr]: 1673, 1548, 1465, 1438, 1404, 1286, 1201, 1180, 1131 cm⁻¹.

HRFABMS: Found 506.25688 calculated for C₂₃H₃₆N₇O₄S 506.25495.

Example 14

N¹,N³-Bis(2-{[5-({[4-({[3-(dimethylamino)propyl]amino}carbonyl)-5-isopropyl-1,3-thiazol-2-yl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]amino}-2-oxoethyl)isophthalamide
N-[3-(Dimethylamino)propyl]-5-isopropyl-2-{{[(1-methyl-4-nitro-1H-pyrrol-2-yl)carbonyl]amino}-1,3-thiazole-4-carboxamide (100 mg, 0.236 mmol; see Example 3, step (iv) above) was dissolved in methanol (25 mL). The solution was cooled to 0°C under N₂ then Pd/C-10% (86 mg) was added. The reaction mixture was hydrogenated at room temperature and atmospheric pressure for 5 h. The catalyst was removed over Kieselguhr and the solvent was then removed under reduced pressure at 50°C. The amine so formed was dissolved in DMF (2 mL, dry), to which

[(3-{{[(carboxymethyl)amino]carbonyl}benzoyl}amino]acetic acid (33 mg, 0.118 mmol; see *J. Med. Chem.* 43, 3257-3266 (2000)) was added followed by HBTU (270 mg, 0.710 mmol) and NMM (300 μ L, dry). An additional amount of DMF (2 mL, dry) was added with stirring at room temperature and the stirring was continued for 72 h. The reaction mixture was diluted with ethyl acetate / methanol (40 mL / 10 mL) and brine with stirring. After extraction the organic layers were collected and the solvents removed under reduced pressure. The product was purified by HPLC and the fractions containing the desired material were freeze-dried. The product was obtained as white solid material (25 mg, 16% yield) with no distinct melting point.

NMR[DMSO- d_6]: 12.03(2H, s); 10.08(2H, s); 9.24(2H, broad); 8.93(2H, t); 8.05(2H, d); 7.94(2H, t); 7.61(1H, t); 7.39(2H, d); 7.25(2H, d); 4.21-4.15(2H, m); 4.06(2H, d); 3.87(6H, s); 3.30(4H, m); 3.07(4H, m); 2.79(6H, d); 1.86(4H, qt); 1.28(12H, d).

IR ν_{\max} [KBr]: 1655, 1548, 1467, 1403, 1288, 1202, 1132 cm^{-1} .

LRLCMS: Found 1029.3 calculated for $\text{C}_{48}\text{H}_{65}\text{N}_{14}\text{O}_8\text{S}_2$ 1029.5.

Example 15

The following compounds were prepared using analogous techniques to those described hereinbefore:

N-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopropyl-1*H*-pyrrol-3-yl]-4-(acetylamino)-1-methyl-1*H*-pyrrole-2-carboxamide; and

N-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]-4-(acetylamino)-1-methyl-1*H*-pyrrole-2-carboxamide.

Example 16

*N*²,*N*⁵-Bis[5-({[4-({[3-(dimethylamino)propyl]amino}carbonyl)-5-isopropyl-1,3-thiazol-2-yl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-1*H*-indole-2,5-dicarboxamide

5 *N*-[3-(Dimethylamino)propyl]-5-isopropyl-2-{[(1-methyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]amino}-1,3-thiazole-4-carboxamide (122 mg, 0.284 mmol; see Example 3, step (iv) above) and Pd/C-10% (110 mg) were suspended in methanol (20 mL) and hydrogenated for 2.5 h at room temperature. Filtration of catalyst over Kieselguhr under N₂, followed by
10 removal of the solvent under reduced pressure gave the amine as an off-white solid, which was used without further purification. Indole-2,5-dicarboxylic acid (29 mg, 0.142 mmol; prepared according to a standard literature procedure (see H. G. Lindwall and G. J. Mantell *J. Org. Chem.* 18, 345-356 (1953))) was dissolved in DMF (0.5 mL, dry), to which was added
15 a solution of HBTU (107 mg, 0.284 mmol) and NMM (0.2 mL, dry) in DMF (1 mL, dry) at room temperature with stirring. The stirring was continued for 30 min. at room temperature. This mixture was added to the amine with stirring at room temperature and stirring was continued at room temperature overnight, followed by purification by HPLC to give the
20 desired material as bis-TFA salt (32 mg, 23% yield) with no distinct melting point.

NMR [DMSO-*d*₆]: 1.3(12H, s, isopropylMe₂); 2.08(1H, m, isopropylCH); 1.85-1.92(4H, m, 2XCH₂); 2.80(6H, s, NMe₂); 2.81(6H, s, NMe₂); 3.07(4H, m, 2xCH₂); 3.23(4H, m, 2xCH₂); 3.93(6H, s, NMe); 7.46(2H, m);
25 7.54(2H, m); 7.85(1H, m); 7.99(2H, m); 8.35(1H, s); 9.37(2H, br, 2xTFA, exch); 10.32(1H, s); 10.52(1H, s); 11.95(1H, s); 12.09(1H, s); 12.15(1H, s).
IR [KBr]: 3415, 2960, 2362, 1654, 1549, 1467, 1398, 1287, 1200, 1134 cm⁻¹.

HRFABMS: found 953.42; calculated for C₄₆H₅₉N₁₃O₆S₂ 953.42.

Example 17*N*²*N*⁵-Bis[1-isopentyl-5-({[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}-carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1*H*-indole-2,5-dicarboxamide

5 1-Isopentyl-*N*-[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]-4-nitro-1*H*-pyrrole-2-carboxamide (120 mg, 0.25 mmol; see Example 30, step (i) below) and Pd/C-10% (110 mg) were suspended in methanol (20 mL) and hydrogenated for 2.5 h at room temperature. Filtration of catalyst over Kieselguhr under N₂, followed by removal of the
10 solvent under reduced pressure gave the amine as an off-white solid, which was used without further purification. Indole-2,5-dicarboxylic acid (26 mg, 0.125 mmol; see Example 16 above) was dissolved in DMF (0.5 mL, dry), to which was added a solution of HBTU (137 mg, 0.36 mmol) and NMM (0.2 mL, dry) in DMF (1 mL, dry) at room temperature with stirring. The
15 stirring was continued for 30 min. at room temperature. This mixture was added to amine with stirring at room temperature and stirring was continued at room temperature overnight, followed by purification by HPLC to give the desired material as bis-TFA salt (15 mg, 12% yield) with no distinct melting point.

20 NMR [DMSO-*d*₆]: 11.95(1H, s); 10.47(1H, s); 10.28(1H, s); 10.02(1H, s); 9.99(1H, s); 9.66(2H, br, TFA); 8.39(1H, s); 8.24(2H, t, unresolved, 2xCONH); 7.88(1H, d, *J*=8.7Hz); 7.49(1H, s); 7.43(2H, s); 7.23(2H, s); 7.14(2H, s); 7.02(2H, s); 4.43(4H, m); 4.06(4H, m); 3.89(6H, s); 3.71(4H, t, *J*=4.6Hz); 3.49(4H, m); 3.31(4H, m); 3.19(8H, m); 1.94(4H, m); 1.66-
25 1.57(6H, m); 0.98(12H, d, *J*=6.5Hz).

IR [KBr]: 1677, 1646, 1533, 1443, 1403, 1198, 1134 cm⁻¹.

LRESMS: Found 1058.6; calculated for C₅₆H₇₆N₁₃O₈ 1058.6

Example 18*N*²,*N*⁵-Bis[5-({[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]-1*H*-indole-2,5-dicarboxamide

5 *N*-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-1-isopentyl-4-nitro-1*H*-pyrrole-2-carboxamide (110 mg, 0.25 mmol; see Example 10, step (i) above) and Pd/C-10% (110mg) were suspended in methanol (20 mL) and hydrogenated for 2.5 h at room temperature. Filtration of catalyst over Kieselguhr under N₂, followed by removal of the
10 solvent under reduced pressure gave the amine as an off-white solid, which was used without further purification. Indole-2,5-dicarboxylic acid (26 mg, 0.125 mmol) was dissolved in DMF (0.5 mL, dry), to which was added a solution of HBTU (137 mg, 0.36 mmol) and NMM (0.2 mL, dry) in DMF (1 mL, dry) at room temperature with stirring. The stirring was continued
15 for 30 min. at room temperature. This mixture was added to amine with stirring at room temperature and stirring was continued at room temperature overnight, followed by purification by HPLC to give the desired material as bis-TFA salt (36mg, 30% yield) with no distinct melting point.

NMR [DMSO-*d*₆]: 11.95(1H, s); 10.47(1H, s); 10.28(1H, s); 10.02(1H, s);
20 9.99(1H, s); 9.39(2H, br, 2xTFA); 8.39(1H, s); 8.22(2H, t, unresolved, 2xCONH); 7.91(1H, d, *J*=8.7Hz); 7.61(1H, d, *J*=8.7Hz); 7.49(1H, s); 7.43(2H, s); 7.23(2H, s); 7.13(2H, s); 7.01(2H, s); 4.43(4H, m); 3.89(6H, s); 3.31(4H, m); 3.14(4H, m); 2.86(12H, d, *J*=3.8Hz); 1.91(4H, m); 1.66-1.55(6H, m); 0.98(12H, d, *J*=6.5Hz).

25 IR [KBr]: 1677, 1647, 1583, 1533, 1443, 1404, 1199, 1135 cm⁻¹.

HRFABMS: Found 974.6; calculated for C₅₂H₇₂N₁₃O₆ 974.5

Example 19

*N*²*N*⁵-Bis[1-isopentyl-5-({[1-methyl-5-({[3-(4-methyl-1-piperazinyl)-propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1*H*-indole-2,5-dicarboxamide

5 1-Isopentyl-*N*-[1-methyl-5-({[3-(4-methyl-1-piperazinyl)propyl]amino]-carbonyl)-1*H*-pyrrol-3-yl]-4-nitro-1*H*-pyrrole-2-carboxamide (122 mg, 0.25 mmol; see Example 29, step (i) below) and Pd/C-10% (110 mg) were suspended in methanol (20 mL) and hydrogenated for 2.5 h at room temperature. Filtration of catalyst over Kieselguhr under N₂, followed by
10 removal of the solvent under reduced pressure gave the amine as an off-white solid, which was used without further purification. Indole-2,5-dicarboxylic acid (26 mg, 0.125 mmol) was dissolved in DMF (0.5 mL, dry), to which was added a solution of HBTU (137 mg, 0.36 mmol) and NMM (0.2 mL, dry) in DMF (1 mL, dry) at room temperature with stirring.
15 The stirring was continued for 30 min. at room temperature. This mixture was added to amine with stirring at room temperature and stirring was continued at room temperature overnight, followed by purification by HPLC to give the desired material as bis-TFA salt (17 mg, 13% yield) with no distinct melting point.

20 NMR [DMSO-*d*₆]: 11.95(1H, s); 10.47(1H, s); 10.27(1H, s); 10.01(1H, s); 9.97(1H, s); 9.60(2H, br, TFA); 8.39(1H, s); 8.13(2H, br); 7.91(1H, d, *J*=8.7Hz); 7.61(1H, d, *J*=8.7Hz); 7.49(1H, s); 7.43(2H, s); 7.21(2H, s); 7.14(2H, s); 6.98(2H, s); 4.43(4H, m); 3.88(6H, s); 3.29(4H, m); 3.20-2.73(20H, m); 1.80(4H, m); 1.66-1.55(6H, m); 0.96(12H, d, *J*=6.5Hz).

25 IR [KBr]: 1676, 1647, 1584, 1533, 1443, 1404, 1200, 1136 cm⁻¹.

HRFABMS: Found 1084.6; calculated for C₅₈H₈₂N₁₅O₆ 1084.6

Example 20

2-({[4-({[4-(Acetylamino)-1-methyl-1*H*-imidazol-2-yl]carbonyl}amino)-1-methyl-1*H*-pyrrol-2-yl]carbonyl}amino)-*N*-[3-(dimethylamino)propyl]-5-isopropyl-1,3-thiazole-4-carboxamide

5 *N*-[3-(Dimethylamino)propyl]-5-isopropyl-2-{{[(1-methyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]amino}-1,3-thiazole-4-carboxamide (105 mg, 0.25 mmol; see Example 3, step (iv) above) and Pd/C-10% (100 mg) were suspended in methanol (20 mL) and hydrogenated for 2.5 h at room temperature. Filtration of catalyst over Kieselguhr under N₂, followed by
10 removal of the solvent under reduced pressure gave the amine as an off-white solid, which was used without further purification. 4-Acetamido-1-methyl-2-imidazolecarboxylic acid sodium salt (120 mg, 0.7 mmol; prepared according to a standard literature procedure (see Tao, Z.; Fujiwara, T.; Saito, I.; Sugiyama, H. *Angew. Chem. Int. Ed. EN* 38(5), 650-653
15 (1999))) was dissolved in DMF (0.5 mL, dry), to which was added a solution of HBTU (137 mg, 0.36 mmol) and NMM (0.2 mL, dry) in DMF (1 mL, dry) at room temperature with stirring. The stirring was continued for 30 min. at room temperature. This mixture was added to the amine with stirring at room temperature and stirring was continued at room temperature
20 overnight, followed by purification by HPLC to give the desired material as bis-TFA salt (30 mg, 22% yield) with no distinct melting point.

NMR [DMSO-*d*₆]: 1.27-1.28(6H, d, *J*=6.8Hz, isopropyl); 1.87(2H, m); 2.02(3H, s); 2.78(6H, s, NMe₂); 3.07(2H, m); 3.32(2H, m); 3.89(3H, s, NMe); 3.95(3H, s, COMe); 4.19(1H, m, isopropyl); 7.44(2H, m); 7.48(1H,
25 m); 7.97(1H, m); 9.3(1H, br); 10.00(1H, s); 10.23(1H, s); 12.03(1H, s).

IR [KBr]: 1667, 1550, 1470, 1288, 1198, 1133 cm⁻¹.

HRFABMS: Found 558.26332 calculated for C₂₅H₃₆N₉O₄S 558.26110.

Example 21

4-(Acetylamino)-N-[1-isopentyl-5-({[1-methyl-5-({[3-(4-methyl-1-piperazinyl)propyl]amino}carbonyl)-1H-pyrrol-3-yl]amino}carbonyl)-1H-pyrrol-3-yl]-1-methyl-1H-pyrrole-2-carboxamide

- 5 *N*-[1-Isopentyl-5-({[1-methyl-5-({[3-(4-methyl-1-piperazinyl)propyl]amino}carbonyl)-1H-pyrrol-3-yl]amino}carbonyl)-1H-pyrrol-3-yl]-1-methyl-4-nitro-1H-pyrrole-2-carboxamide (186 mg, 0.305 mmol; see Example 29, step (ii) below) was dissolved in methanol at 0°C under N₂ with stirring to which Pd/C-10% (136 mg) was added. The reaction mixture
10 was hydrogenated for 3 h at room temperature and atmospheric pressure with stirring. The catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure. The amine so formed was dissolved in dichloromethane (10 mL) and this solution was divided in half. Acetyl chloride (12 mg, 12 µL, 0.153 mmol) was added dropwise with stirring at
15 room temperature to the first half. The stirring was continued at room temperature overnight. The solvent was removed under reduced pressure and the crude product formed was dissolved in small amount of acetonitrile/water containing 0.1% TFA and purified by HPLC. Fractions containing the pure material were combined and freeze-dried to give off-
20 white solid (45.5 mg, 40% yield) with no distinct melting point.

NMR [DMSO-d₆]: 9.87(1H, s); 9.86(1H, s); 9.79(1H, s); 8.09(1H, t, unresolved); 7.26(1H, d, J=1.7Hz); 7.14(1H, d, J=1.7Hz); 7.13(1H, d, J=1.7Hz); 7.01(1H, d, J=1.7Hz); 6.91(1H, d, J=1.7Hz); 6.86(1H, d, J=1.7Hz); 4.31(2H, t, J=6.9Hz); 3.83(3H, s); 3.81(3H, s); 3.24(2H, q, J=6.4Hz); 3.10-2.60(8H, br, & 3H, s); 1.97(3H, s); 1.77(2H, m); 1.58-1.46(3H, m); 0.90(6H, d, J=6.4Hz).

IR [KBr]: 1675, 1647, 1584, 1535, 1459, 1436, 1402, 1197, 1132 cm⁻¹.

HRFABMS: Found 622.38301; calculated for C₃₂H₄₈N₉O₄ 622.38293.

Example 22

N-[1-Isopentyl-5-({[1-methyl-5-({[3-(4-methyl-1-piperazinyl)propyl]-amino}carbonyl)-1H-pyrrol-3-yl]amino}carbonyl)-1H-pyrrol-3-yl]-4-[(3-methoxybenzoyl)amino]-1-methyl-1H-pyrrole-2-carboxamide

- 5 To the second half of the amine solution (see Example 21 above) was added *m*-methoxybenzoyl chloride (26 mg, 26 μ L, 0.153 mmol) was added dropwise at room temperature with stirring. The stirring was continued overnight at room temperature. The solvent was removed under reduced pressure and the crude product was purified by HPLC. The product was
10 obtained as light brown solid (60.8 mg, 48% yield).

NMR [DMSO- d_6]: 10.28(1H, s); 9.95(1H, s); 9.88(1H, s); 8.09(1H, t, unresolved); 7.53-7.47(2H, m); 7.43(1H, t, $J=7.8$ Hz); 7.32(1H, d, $J=1.7$ Hz); 7.29(1H, d, $J=1.7$ Hz); 7.15-7.11(3H, m); 7.02(1H, d, $J=1.7$ Hz); 6.92(1H, d, $J=1.7$ Hz); 4.32(2H, t, $J=6.6$ Hz); 3.88(3H, s); 3.83(3H, s); 3.81(3H, s);
15 3.24(2H, q, $J=6.0$ Hz); 3.10-2.40(8H, br, & 3H, s); 1.77(2H, br); 1.59-1.47(3H, m); 0.91(6H, d, $J=6.4$ Hz).

IR [KBr]: 1677, 1645, 1583, 1535, 1462, 1435, 1402, 1263, 1197, 1132 cm^{-1} .

HRFABMS: Found 714.40764; calculated for $\text{C}_{38}\text{H}_{52}\text{N}_9\text{O}_5$ 714.40914.

20

Example 23

N-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]-4-({[5-(formylamino)-2-methyl-3-thienyl]carbonyl}amino)-1-isopentyl-1H-pyrrole-2-carboxamide

25

(i) Ethyl 2-methyl-4,5-dihydro-3-thiophenecarboxylate

A standard procedure was employed (see Pallab Chatterjee, Patrick J. Murphy, Rosanna Pepe and Michael Shaw *J. Chem. Soc., Perkin Trans. 1* 17, 2403-2406 (1994)) to give the product as colourless oil (96% yield).

(ii) Ethyl 2-methyl-3-thiophenecarboxylate

A standard procedure was employed (see Pallab Chatterjee, Patrick J. Murphy, Rosanna Pepe and Michael Shaw *J. Chem. Soc., Perkin Trans. 1* 17, 2403-2406 (1994)) to give the product as colourless oil (82% yield).

5

(iii) 2-Methyl-3-thiophenecarboxylic acid

Ethyl 2-methyl-3-thiophenecarboxylate (3.42g, 20.09 mmol; see step (ii) above) was dissolved in ethanol (5 mL) to which was added a solution of sodium hydroxide (3.2145 mg, 80.362 mmol) in water (10 mL). The reaction mixture was heated under reflux for 2 h. The volume was reduced to half under reduced pressure at 40°C and the residue was cooled with ice water. Dilute HCl was added dropwise with stirring until pH 2. The white solid material was filtered off, washed with water and dried under reduced pressure at 50°C to give the required material (2.1850 mg, 77% yield), m.p. 115-117°C, lit. m.p. 116-117°C (see D.W. Knight, A.P. Nott *J. Chem. Soc., Perkin Trans. 1* 791-794 (1983)).

10
15

(iv) 2-Methyl-5-nitro-3-thiophenecarboxylic acid

A mixture of concentrated nitric acid (10 mL, sp.gr. 1.42) and concentrated sulphuric acid (6 mL) was mechanically stirred in a round-bottomed flask and cooled to (-10°C) by a dry ice methanol bath. The temperature was kept below (-5°C) while 2-methyl-3-thiophenecarboxylic acid (996 mg, 7.006 mmol; see step (iii) above) was added in small portions. The reaction mixture was stirred at the same temperature for 15 min. and then was poured over ice water. The solid material that precipitated was filtered off, washed with water and dried to give light brown solid (883 mg; 67% yield), m.p. 177-180°C.

20
25

NMR [DMSO-d₆]: 13.35(1H, br); 8.11(1H, s); 2.76(3H, s).

IR [KBr]: 1706, 1543, 1514, 1457, 1335, 1257cm⁻¹.

30 HREIMS: Found 186.99431; calculated for C₆H₅NO₄S 186.99393.

(v) *N*-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-1-isopentyl-4-{[(2-methyl-5-nitro-3-thienyl)carbonyl]amino}-1*H*-pyrrole-2-carboxamide

5 *N*-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-1-isopentyl-4-nitro-1*H*-pyrrole-2-carboxamide (204mg, 0.463mmol; see Example 10, step (i) above) was dissolved in methanol (25 mL). The reaction mixture was cooled to 0°C and then Pd/C-10% (121 mg) was added portion-wise under N₂ with stirring. The reaction mixture was
10 hydrogenated for 5h at room temperature and atmospheric pressure. The catalyst was removed over Kieselguhr and the solvent was then removed under reduced pressure. 2-Methyl-5-nitro-3-thiophenecarboxylic acid (87 mg, 0.463 mmol; see step (iv) above) was dissolved in thionyl chloride (2mL) then heated until reflux for 5 h. The excess thionyl chloride was
15 removed under reduced pressure to give the acid chloride, which was dissolved in DCM (5 mL). The amine was dissolved in DCM (5 mL) to which *N*-methyl morpholine (0.1 mL) was added followed by the acid chloride, which was added dropwise at room temperature with stirring. The stirring was continued at room temperature overnight. The volatile material
20 was removed under reduced pressure and the crude product was then purified by column chromatography using 2.5/2.5/0.6 mL ethyl acetate/methanol/TEA) R_F=0.5. The product was obtained as amorphous yellow material (237 mg, 90% yield) with no distinct melting point.

NMR [DMSO-d₆]: 10.37(1H, s); 9.94(1H, s); 8.58(1H, s); 8.12(1H, t, J=5.6Hz); 7.35(1H, d, J=1.7Hz); 7.18(1H, d, J=1.7Hz); 7.14(1H, d, J=1.7Hz); 7.01(1H, d, J=1.7Hz); 6.86(1H, d, J=1.7Hz); 4.34(1H, t, J=6.9Hz); 3.80(3H, s); 3.22(2H, q, J=6.5Hz); 2.77(3H, s); 2.64(2H, m); 2.44(6H, s); 1.73(2H, quintet, J=6.7Hz); 1.59-1.45(3H, m); 0.90(6H, d, J=6.4Hz).

IR [KBr]: 1675, 1650, 1583, 1537, 1465, 1437, 1403, 1263, 1202, 1133 cm⁻¹.

HRFABMS: Found: 572.26551; calculated for C₂₇H₃₇N₇O₅S 572.26580.

- 5 (vi) *N*-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-4-({[5-(formylamino)-2-methyl-3-thienyl]carbonyl}amino)-1-isopentyl-1*H*-pyrrole-2-carboxamide

N-[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-1-isopentyl-4-{{(2-methyl-5-nitro-3-thienyl)carbonyl]amino}-1*H*-

- 10 pyrrole-2-carboxamide (110 mg, 0.193 mmol; see step (v) above) was dissolved in ethanol (20 mL) and cooled to 0°C. Pd/C-10% (120 mg) was added under N₂ with stirring and the reaction was hydrogenated at room temperature and atmospheric pressure for 3 h. The catalyst was removed over Kieselguhr then ethyl formate (20 mL) was added to the ethanolic
15 solution. The reaction mixture was heated under reflux for 3 days. The volatile material was then removed under reduced pressure and the crude product was purified by HPLC. The product was obtained as solid material with no distinct melting point (23 mg, 17.4% yield).

NMR [DMSO-d₆]: 11.32(1H, s); 9.94(1H, s); 9.88(1H, s); 9.24(1H, br, TFA); 8.29(1H, s); 8.14(1H, t, J=5.9Hz); 7.29(1H, d, J=1.7Hz); 6.95(1H, s); 6.93(1H, d, J=1.7Hz); 4.32(1H, t, J=7.0Hz); 3.81(3H, s); 3.25(2H, q, J=5.1Hz); 3.06(2H, m); 2.79(6H, d, J=3.5Hz); 2.5(3H, s); 1.83(2H, quintet, J=6.7Hz); 1.57-1.49(3H, m); 0.90(6H, d, J=6.4Hz).

IR [KBr]: 1671, 1651, 1582, 1537, 1464, 1403, 1200, 1179, 1132 cm⁻¹.

- 25 HRFABMS: Found: 570.28565; calculated for C₂₈H₄₀N₇O₄S 570.28625.

Example 24

N-[5-({[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]-5-isopropyl-2-[(3-methoxybenzoyl)amino]-1,3-thiazole-4-carboxamide

5

(i) Methyl 5-isopropyl-2-[(3-methoxybenzoyl)amino]-1,3-thiazole-4-carboxylate

Methyl 2-amino-5-isopropyl-1,3-thiazole-4-carboxylate (383 mg, 1.913 mmol; see Example 3, step (i) above) was dissolved in DCM (10 mL) at room temperature with stirring. N-methyl morpholine (0.3mL) was added followed by 3-methoxybenzoyl chloride (326 mg, 1.913 mmol) at room temperature with stirring. The reaction mixture was heated until reflux for 10 min. and then was left stirring at room temperature overnight. The solvent was removed under reduced pressure and the crude product was purified by column chromatography using silica gel and 1:2 ethyl acetate/petroleum ether. Fractions containing the required material ($R_F=0.4$) were collected and the solvents were removed under reduced pressure to give the product as white solid material (530 mg, 83% yield), m.p.60-63°C.

NMR [$CDCl_3$]: 9.89(1H, s); 7.45-7.38(3H, m); 7.16-7.13(1H, m); 4.16(1H, heptet, $J=6.8$ Hz); 3.86(3H, s); 3.85(3H, s); 1.40(6H, d, $J=6.8$ Hz).

IR [KBr]: 2958, 1720, 1669, 1547, 1463, 1298, 1208, 1045, 822, 743 cm^{-1} .

HRFABMS: Found: 335.10620; calculated for $C_{16}H_{19}N_2O_4S$ 335.10655.

(ii) 5-Isopropyl-2-[(3-methoxybenzoyl)amino]-1,3-thiazole-4-carboxylic acid

Methyl 5-isopropyl-2-[(3-methoxybenzoyl)amino]-1,3-thiazole-4-carboxylate (330 mg, 0.998 mmol; see step (i) above) was dissolved in ethanolic KOH (561 mg, in 10 mL ethanol; 10 mmol). The reaction mixture was heated under reflux for 3 h, and then it was cooled to 0°C.

HCl_(conc.) was added dropwise with stirring at 0°C and the resulting solid was filtered off, washed with water and dried under reduced pressure at 45°C overnight. The product was obtained as white solid material (290 mg, 91% yield), m.p.>230°C [sublimation].

5 NMR [DMSO-d₆]: 7.69(2H, m); 7.44(1H, t, J=3.9Hz); 7.19(1H, m); 4.06(1H, heptet, J=6.8Hz); 3.84(3H, s); 1.31(6H, d, J=6.8Hz).

IR [KBr]: 2963, 1672, 1555, 1463, 1306, 1221, 1043, 936, 820, 736 cm⁻¹.

HRFABMS: Found: 321.09056; calculated for C₁₅H₁₇N₂O₄S 321.09090.

10 (iii) N-[5-({[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]-5-isopropyl-2-[(3-methoxybenzoyl)amino]-1,3-thiazole-4-carboxamide

N-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]-1-methyl-4-nitro-1H-pyrrole-2-carboxamide (113 mg, 0.300 mmol; see
15 *Tetrahedron* 56, 5225-5239, (2000)) was suspended in methanol (25 mL) at 0°C with stirring under N₂. Pd/C-10% (61 mg) was added with stirring under N₂ and then the reaction mixture was hydrogenated at room temperature and atmospheric pressure for 3 h. The catalyst was removed over Kieselguhr and the solvent was then removed under reduced pressure
20 to give the amine, which was used in the coupling reaction without further purification. 5-Isopropyl-2-[(3-methoxybenzoyl)amino]-1,3-thiazole-4-carboxylic acid (96 mg, 3.00 mmol; see step (ii) above) was dissolved in thionyl chloride (2 mL) then heated under reflux for 3 h. The excess thionyl chloride was removed under reduced pressure and the acid chloride so
25 formed was dissolved in DCM (5 mL). The amine was dissolved in DCM (5 mL), to which N-methyl morpholine (0.2 mL) was added. The acid chloride solution was added dropwise at room temperature with stirring and then the stirring was continued overnight at room temperature. The volatile material was removed under reduced pressure and the crude product was
30 purified by HPLC. Fractions containing the required material were

collected and freeze-dried to give the product as pale yellow solid (73 mg, 32% yield) with no distinct melting point.

NMR [DMSO- d_6]: 12.63(1H, s); 9.88(1H, s); 9.55(1H, s); 9.29(1H, br); 8.14(1H, t, 5.8Hz); 7.69(1H, s); 7.67(1H, s); 7.48(1H, t, $J=5.9$ Hz); 7.26(1H, d, 1.7Hz); 7.23(1H, m); 7.18(1H, d, $J=1.7$ Hz); 7.13(1H, d, $J=1.7$ Hz); 6.94(1H, d, 1.7Hz); 4.19(1H, heptet, $J=6.8$ Hz); 3.86(6H, s); 3.82(3H, s); 3.24(2H, q, $J=6.7$ Hz); 3.07(2H, m); 2.79(6H, d, $J=3.8$ Hz); 1.86(2H, quintet, $J=6.7$ Hz); 1.33(6H, d, $J=6.8$ Hz).

IR [KBr]: 1660, 1547, 1466, 1436, 1403, 1286, 1199, 1137, 1043 cm^{-1} .

HRFABMS: Found: 649.29379; calculated for $\text{C}_{32}\text{H}_{41}\text{N}_8\text{O}_5\text{S}$ 649.29206.

Example 25

N-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-4-{{[(5-{{[(9,10-dioxo-9,10-dihydro-2-anthracenyl)carbonyl]amino}-2-

15 methyl-3-thienyl)carbonyl]amino}-1-isopentyl-1*H*-pyrrole-2-carboxamide

N-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-1-isopentyl-4-{{[(2-methyl-5-nitro-3-thienyl)carbonyl]amino}-1*H*-

pyrrole-2-carboxamide (127 mg, 0.223 mmol; see Example 23, step (v) above) was dissolved in methanol (25mL) at 0°C under N_2 with stirring.

20 Pd/C-10% (114 mg) was added and the reaction mixture was hydrogenated at room temperature and atmospheric pressure for 3.5 h. The catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure to give the amine, which was used in the subsequent coupling reaction. 9,10-Dioxo-9,10-dihydro-2-anthracenecarboxylic acid (Aldrich)

25 (56 mg, 0.223 mmol) was dissolved in DMF (1 mL, dry), to which HBTU (127 mg, 0.335 mmol) and *N*-methyl morpholine (0.2 mL, dry) were added with stirring under N_2 . The amine was dissolved in DMF (1mL, dry) and was added dropwise to the reaction mixture. The reaction mixture was left stirring at room temperature overnight. The reaction mixture was then

30 diluted with a mixture made of ethyl acetate/methanol 9:1 (50 mL), after

which it was extracted with water. The organic layer was dried over MgSO_4 , filtered and the solvents removed under reduced pressure. The crude product obtained was purified by HPLC to give the pure material as brown solid with no distinct melting point (33 mg, 19% yield).

5 NMR [DMSO-d_6]: 21.04(1H, s); 10.04(1H, s); 9.91(1H, s); 9.23(1H, br, TFA); 8.83(1H, d, $J=1.7\text{Hz}$); 8.50(1H, dd, $J=1.8\text{Hz}$ & $J=8.1\text{Hz}$); 8.42(1H, d, $J=8.1\text{Hz}$); 8.29-8.25(2H, m); 8.15(1H, t, $J=5.9\text{Hz}$); 8.00-7.97(2H, m); 7.33(1H, d, $J=1.7\text{Hz}$); 7.21(1H, s); 7.17(1H, d, $J=1.7\text{Hz}$); 7.02(1H, d, $J=1.7\text{Hz}$); 6.94(1H, d, $J=1.7\text{Hz}$); 4.33(2H, t, $J=7.1\text{Hz}$); 3.82(3H, s);
10 3.24(2H, q, $J=5.1\text{Hz}$); 3.08(2H, m); 2.79(6H, d, $J=3.5\text{Hz}$); 2.59(3H, s); 1.84(2H, quintet, $J=6.7\text{Hz}$); 1.60-1.50(3H, m); 0.91(6H, d, $J=6.4\text{Hz}$).
IR [KBr]: 1674, 1653, 1585, 1539, 1464, 1404, 1487, 1200, 1177, 1135 cm^{-1} .

HRFABMS: Found: 776.32284; calculated for $\text{C}_{42}\text{H}_{46}\text{N}_7\text{O}_6\text{S}$ 776.32303.

15

Example 26

N-[1-(Cyclopropylmethyl)-5-({[5-({[3-(dimethylamino)propyl]amino}-carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-4-(formylamino)-1-methyl-1*H*-pyrrole-2-carboxamide

20

(i) Ethyl 1-(cyclopropylmethyl)-4-nitro-1*H*-pyrrole-2-carboxylate

Ethyl 4-nitro-1*H*-pyrrole-2-carboxylate (1.042 g, 5.626 mmol; see Example 1, step (i) above) was dissolved in DMF (20 mL, dry) to which potassium metal (0.435 g, 11.124 mmol) was added. The reaction mixture was heated
25 to 100°C with stirring and it was left at that temperature for 1 h. The reaction mixture was cooled to 50-60°C then (chloromethyl)cyclopropane (1.000 g, 11.044 mmol) [purchased from Aldrich] and KI (1.349 g, 8.127 mmol) were added and the reaction mixture was heated again with stirring to 80°C for 5 h, after which time it was left stirring overnight at
30 room temperature. The reaction mixture was diluted with brine (50 mL)

and extracted with ethyl acetate (3x100 mL). The combined organic extracts were collected and the solvent was removed under reduced pressure. The crude product obtained (containing some DMF) was purified by flash chromatography using silica gel and 1/10 ethyl acetate/*n*-hexane.

- 5 The product as white microcrystalline solid [$R_F=0.2$] was obtained (1.070 g, 80% yield), m.p. 65-66°C.

NMR [DMSO- d_6]: 8.34(1H, d, $J=1.3$ Hz); 7.35(1H, d, $J=1.3$ Hz); 4.30-4.24(2H, q, $J=7.1$ Hz); 4.21(2H, d, $J=7.2$ Hz); 1.34-1.23(2H, t, $J=7.1$ Hz & 1H, m); 0.53-0.45(2H, m); 0.42-0.38(2H, m).

- 10 IR [KBr]: 3142, 2996, 2921, 1704, 1502, 1310, 1270, 1225, 1184, 1091 cm^{-1} .

HREIMS: Found 238.09585; calculated for $\text{C}_{11}\text{H}_{14}\text{N}_2\text{O}_4$ 238.09536.

(ii) 1-(Cyclopropylmethyl)-4-nitro-1*H*-pyrrole-2-carboxylic acid

- 15 Ethyl 1-(cyclopropylmethyl)-4-nitro-1*H*-pyrrole-2-carboxylate (482 mg, 2.023 mmol; see step (i) above) was dissolved in ethanol (4 mL) to which a solution of NaOH (393 mg, 9.825 mmol) in water (10 mL) was added. The reaction mixture was heated under reflux for 2 h then cooled to 0°C. HCl conc. was added dropwise with stirring until the solution reached pH 2. The
20 white solid material, which precipitated was filtered off, washed with water and dried under reduced pressure at 60°C overnight. The product was obtained as white solid material (382 mg, 90% yield), m.p. 215-217°C.

NMR [DMSO- d_6]: 13.14(1H, br); 8.29(1H, d, $J=2.4$ Hz); 7.29(1H, d, $J=2.1$ Hz); 4.20(2H, d, $J=2.2$ Hz); 1.34-1.27(1H, m); 0.52-0.38(4H, m).

- 25 IR [KBr]: 3115, 3005, 2922, 2854, 1680, 1540, 1517, 1481, 1420, 1314 cm^{-1} .

HREIMS: Found 210.06469; calculated for $\text{C}_9\text{H}_{10}\text{N}_2\text{O}_4$ 210.06406.

(iii) 1-(Cyclopropylmethyl)-N-[5-({[3-(dimethylamino)propyl]amino}-carbonyl)-1-methyl-1H-pyrrol-3-yl]-4-nitro-1H-pyrrole-2-carboxamide
N-[3-(Dimethylamino)propyl]-1-methyl-4-nitro-1*H*-pyrrole-2-carboxamide
(100 mg, 0.393 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) was
5 dissolved in methanol (25 mL) to which Pd/C-10% (75 mg) was added at
0°C under N₂ with stirring. The reaction mixture was hydrogenated for 2 h
at room temperature and atmospheric pressure. The catalyst was removed
over Kieselguhr and the solvent was removed under reduced pressure. The
amine so formed was dissolved in DMF (2 mL, dry) to which was added
10 1-(cyclopropylmethyl)-4-nitro-1*H*-pyrrole-2-carboxylic acid (83 mg,
0.393 mmol; see step (ii) above), HBTU (298 mg, 0.786 mmol), NMM
(200 µL, dry) with stirring at room temperature. The stirring was continued
for 48 h. The reaction mixture was diluted with ethyl acetate (50 mL) and
extracted with NaHCO₃ (25 mL, saturated). The water layer was extracted
15 again with ethyl acetate (50 mL). The combined organic extract was dried
over MgSO₄, filtered and the solvent removed under reduced pressure to
give the crude product as pale yellow oil. Purification by flash column
chromatography over silica gel using 1/2/0.1 methanol/ethyl acetate/TEA
gave the required product as glassy yellow material [*R*_F=0.5] (130 mg, 80%
20 yield) with no distinct melting point.

NMR [DMSO-*d*₆]: 10.26(1H, s); 8.24(1H, d, *J*=1.9Hz); 8.10(1H, t,
J=5.6Hz); 7.57(1H, d, *J*=1.9Hz); 7.21(1H, d, *J*=1.9Hz); 6.83(1H, d,
J=1.9Hz); 4.27(2H, d, *J*=7.2Hz); 3.81(3H, s); 3.21-3.16(2H, q, *J*=6.9Hz);
2.28(2H, t, *J*=7.1Hz); 2.17(6H, s); 1.65-1.59(2H, quintet, *J*=7.1Hz); 1.34-
25 1.29(1H, m); 0.51-0.38(4H, m).

IR [KBr]: 3283, 3126, 2944, 1642, 1573, 1532, 1504, 1463, 1427, 1389,
1309, 1233 cm⁻¹.

HRFABMS: Found 417.22460; calculated for C₂₀H₂₉N₆O₄ 417.22503.

(iv) *N*-[1-(Cyclopropylmethyl)-5-({[5-({[3-(dimethylamino)propyl]amino}-carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1-methyl-4-nitro-1*H*-pyrrole-2-carboxamide

1-(Cyclopropylmethyl)-*N*-[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-4-nitro-1*H*-pyrrole-2-carboxamide (120 mg, 0.288 mmol; see step (iii) above) was dissolved in methanol (25 mL) to which Pd/C-10% (97 mg) was added at 0°C under N₂ with stirring. The reaction mixture was hydrogenated for 4 h at room temperature and atmospheric pressure. The catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure to give the amine, which was used in the next step without any purification. The amine so formed was dissolved in DMF (2 mL, dry) to which 1-methyl-4-nitro-1*H*-pyrrole-2-carboxylic acid (57 mg, 0.288 mmol; see *Tetrahedron* 56, 5225-5239 (2000)), HBTU (218 mg, 0.576 mmol), and NMM (200 µL, dry) were added at room temperature with stirring. The stirring was continued at room temperature overnight. The reaction mixture was diluted with ethyl acetate (50 mL) and extracted with NaHCO₃ (25 mL, sat.). The water layer was extracted again with ethyl acetate (50 mL). The combined organic extract was dried over MgSO₄, filtered and the solvent was removed under reduced pressure. The crude product obtained was purified by column chromatography using silica gel and 1/2/0.1 methanol/ethyl acetate/TEA. Fractions containing the pure product [*R*_F=0.4] were collected and the solvents were removed under reduced pressure at 50°C to give the required product as glassy yellow solid (110 mg, 71% yield) with no distinct melting point.

NMR [DMSO-*d*₆]: 10.27(1H, s); 9.94(1H, s); 8.18(1H, d, *J*=1.7Hz); 8.06(1H, t, *J*=5.2Hz); 7.59(1H, d, *J*=1.9Hz); 7.36(1H, d, 1.9Hz); 7.18(1H, d, *J*=1.9Hz); 7.02(1H, d, *J*=1.9Hz); 6.83(1H, d, *J*=1.9Hz); 4.19(2H, d, *J*=7.0Hz); 3.97(3H, s); 3.80(3H, s); 3.21-3.16(2H, q, *J*=6.9Hz); 2.26(2H, t,

J=7.1Hz); 2.15(6H, s); 1.65-1.58(2H, quintet, J=7.1Hz); 1.26-1.13(1H, m); 0.47-0.42(2H, m); 0.33-0.31(2H, m).

IR [KBr]: 3286, 3126, 2938, 1646, 1580, 1531, 1462, 1439, 1400, 1309, 1248 cm⁻¹.

5 HRFABMS: Found 539.27295; calculated for C₂₆H₃₅N₈O₅ 539.27304.

(v) N-[1-(Cyclopropylmethyl)-5-({[5-({[3-(dimethylamino)propyl]amino}-carbonyl)-1-methyl-1H-pyrrol-3-yl]amino}carbonyl)-1H-pyrrol-3-yl]-4-(formylamino)-1-methyl-1H-pyrrole-2-carboxamide

10 N-[1-(Cyclopropylmethyl)-5-({[5-({[3-(dimethylamino)propyl]amino}-carbonyl)-1-methyl-1H-pyrrol-3-yl]amino}carbonyl)-1H-pyrrol-3-yl]-1-methyl-4-nitro-1H-pyrrole-2-carboxamide (100mg, 0.186 mmol; see step (iv) above) was dissolved in ethanol (25 mL) to which Pd/C-10% (78 mg) was added at 0°C under N₂ with stirring. The reaction mixture was
15 hydrogenated for 3 h at room temperature and atmospheric pressure. The catalyst was removed over Kieselguhr. Ethyl formate was added to the ethanolic solution of the amine and the reaction mixture was heated under reflux for 48 h. Ethanol and excess ethyl formate were removed under reduced pressure and the crude product was purified by HPLC. The product
20 was obtained as pale yellow solid (52 mg, 42% yield) after freeze drying with no distinct melting point.

NMR [DMSO-d₆]: 10.04(1H, s); 9.91(1H, s); 9.90(1H, s); 9.28(1H, br, TFA); 8.16-8.13(2H, t&d); 7.31(1H, d, J=1.7Hz); 7.19(1H, d, J=1.7Hz); 7.16(1H, d, J=1.7Hz); 7.05(1H, d, J=1.7Hz); 6.95(1H, d, 1.7Hz); 6.93(1H, d, J=1.7Hz); 4.18(2H, d, J=7.0Hz); 3.85(3H, s); 3.81(3H, s); 3.26-3.22(2H, q, J=6.3Hz); 3.07(2H, m); 2.79(2H, d, J=4.3Hz); 1.87-1.82(2H, quintet, J=6.6Hz); 1.23(1H, m); 0.47-0.42(2H, m); 0.33-0.29(2H, m).

IR[KBr]: 3410, 3294, 1674, 1649, 1582, 1533, 1464, 1438, 1403, 1201, 1132 cm⁻¹.

30 HRFABMS: Found 537.29360; calculated for C₂₇H₃₇N₈O₄ 537.29378.

Example 27

1-Cyclopentyl-N-[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]-4-({[4-(formylamino)-1-methyl-1H-pyrrol-2-yl]-carbonyl}amino)-1H-pyrrole-2-carboxamide

(i) Ethyl 1-cyclopentyl-4-nitro-1H-pyrrole-2-carboxylate

Ethyl 4-nitro-1H-pyrrole-2-carboxylate (1.006 g, 5.432 mmol; see Example 1, step (i) above) was dissolved in DMF (20 mL, dry) to which was added potassium metal (0.492 g, 12.582 mmol) then it was heated to 80°C and then it was left at that temperature for 1 h with stirring. The reaction mixture was cooled to 50-60°C after which bromocyclopentane (1 mL, 1.390 g, 9.326 mmol) was added followed by KI (1.548 g, 9.326 mmol). The reaction mixture was heated again for 5 h with stirring at 80°C, after which time it was left stirring at room temperature under N₂, overnight. The reaction mixture was diluted with brine under N₂, at room temperature, and then it was extracted with ethyl acetate (3X100 mL). The combined organic extract was dried over MgSO₄, filtered and the solvent removed under reduced pressure to give the crude product (containing some DMF). The product was purified using column chromatography [silica gel, 1/10 ethyl acetate/n-hexane (R_F=0.24)]. Fractions containing the pure material were collected and the solvent removed to give pale yellow oil (440 mg, 32% yield).

NMR [CDCl₃]: 7.77(1H, d, J=2.0Hz); 7.45(1H, d, J=2.0Hz); 5.57-5.50(1H, quintet, J=6.6Hz); 4.33(2H, q, J=7.1Hz); 2.30-2.24(2H, m); 1.98-1.75(6H, m); 1.37(3H, t, J=7.1Hz).

IR[KBr]: 2965, 2875, 1717, 1534, 1508, 1426, 1335, 1294, 1222, 1187, 1084 cm⁻¹.

HRFABMS: Found 253.11824; calculated for C₁₂H₁₇N₂O₄ 253.11883.

(ii) 1-Cyclopentyl-4-nitro-1H-pyrrole-2-carboxylic acid

Ethyl 1-cyclopentyl-4-nitro-1H-pyrrole-2-carboxylate (430 mg, 1.706 mmol; see step (i) above) was dissolved in ethanol (4 mL) to which a solution of NaOH (408 mg, 10.20 mmol) in water (10 mL) was added. The reaction mixture was heated under reflux for 2 h, and then it was cooled to 0°C. Hydrochloric acid (conc.) was added dropwise with stirring until pH 2. The solid was filtered off, washed with water and dried at 45°C for 48 h to give white solid material (340 mg, 89% yield). Mp 195-198°C.

NMR [DMSO-d₆]: 13.13(1H, br); 8.27(1H, d, J=2.0Hz); 7.29(1H, d, J=2.0Hz); 5.45(1H, quintet, J=7.1Hz); 2.16-2.07(2H, m); 1.88-1.78(4H, m); 1.65-1.61(2H, m).

IR [KBr]: 3152, 2964, 2880, 1683, 1508, 1430, 1331, 1292, 1078, 911 cm⁻¹.

HREIMS: Found 224.07926; calculated for C₁₀H₁₂N₂O₄ 224.07971.

(iii) 1-Cyclopentyl-N-[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]-4-nitro-1H-pyrrole-2-carboxamide

N-[3-(Dimethylamino)propyl]-1-methyl-4-nitro-1H-pyrrole-2-carboxamide (100 mg, 0.393 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) was dissolved in methanol (25 mL) to which Pd/C-10% (90 mg) was added at 0°C under N₂ with stirring. The reaction mixture was hydrogenated for 3 h at room temperature and atmospheric pressure. The catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure. The amine so formed was dissolved in DMF (2 mL, dry) to which 1-(cyclopentyl-4-nitro-1H-pyrrole-2-carboxylic acid (88 mg, 0.393 mmol; see step (ii) above) was added followed by HBTU (298 mg, 0.786 mmol) and NMM (200 µL, dry) with stirring at room temperature. The reaction mixture was diluted with ethyl acetate (50 mL) and extracted with NaHCO₃(sat.) (25 mL). The water layer was extracted again with ethyl acetate (50 mL). The combined organic extract was dried over MgSO₄, filtered and the solvent removed under reduced pressure to give the crude

product which was purified by column chromatography using silica gel and 1/2/0.1 methanol/ethyl acetate/TEA. The product was obtained as yellow oil $R_F=0.45$ (157 mg, 93% yield).

NMR [DMSO- d_6]: 10.27(1H, s); 8.24(1H, d, $J=1.7$ Hz); 8.11(1H, t, $J=5.2$ Hz); 7.47(1H, s); 7.20(1H, d, $J=1.7$ Hz); 6.83(1H, d, $J=1.7$ Hz); 5.48(1H, quintet, $J=7.4$ Hz); 3.80(3H, s); 3.18(2H, q, $J=6.6$ Hz); 2.66(2H, m); 2.30(2H, t, $J=7.1$ Hz); 2.14(6H, s); 2.12(2H, m); 1.82(4H, m); 1.62(2H, quintet, $J=7.1$ Hz).

IR [KBr]: 1644, 1575, 1534, 1504, 1437, 1401, 1313, 1287, 748 cm^{-1} .

HRFABMS: Found 431.24204; calculated for $C_{21}H_{31}N_6O_4$ 431.24068.

(iv) 1-Cyclopentyl-N-[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]-4-{{[(1-methyl-4-nitro-1H-pyrrol-2-yl)carbonyl]-amino}-1H-pyrrole-2-carboxamide

1-Cyclopentyl-N-[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]-4-nitro-1H-pyrrole-2-carboxamide (157 mg, 0.365 mmol; see step (iii) above) was dissolved in methanol (25 mL) at 0°C under N_2 with stirring, to which Pd/C-10% (80 mg) was added. The reaction mixture was hydrogenated for 2 h at room temperature and atmospheric pressure. The catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure. The amine so formed was dissolved in DMF (2 mL, dry) to which 1-methyl-4-nitro-1H-pyrrole-2-carboxylic acid (62 mg, 0.365 mmol; see *Tetrahedron* 56, 5225-5239 (2000)), HBTU (277 mg, 0.730 mmol) and NMM (200 μ L, dry) were added with stirring at room temperature. The stirring was continued overnight. The reaction mixture was diluted with ethyl acetate (50 mL) and extracted with $NaHCO_3$ (sat) (25 mL). The water layer was extracted again with ethyl acetate (50 mL). The combined organic extract was dried over $MgSO_4$, filtered and the solvent removed under reduced pressure to give the crude product, which was purified by column chromatography using silica gel and

1/2/0.1 methanol/ethyl acetate/TEA, $R_F=0.40$. The product was obtained as glassy solid material with no distinct melting point (110 mg, 55% yield).

NMR [DMSO- d_6]: 10.28(1H, s); 9.96(1H, s); 8.18(1H, d, $J=1.8$ Hz); 8.07(1H, t, $J=5.2$ Hz); 7.58(1H, d, $J=1.8$ Hz); 7.41(1H, d, $J=1.8$ Hz); 7.18(1H, d, $J=1.8$ Hz); 6.93(1H, d, $J=1.8$ Hz); 6.85(1H, d, $J=1.8$ Hz); 5.53(1H, quintet, $J=7.3$ Hz); 3.97(3H, s); 3.80(3H, s); 3.20(2H, m); 2.83(2H, m); 2.42(2H, t, $J=7.2$ Hz); 2.27(6H, s); 2.09(2H, m); 1.79-1.64(6H, m).

IR [KBr]: 1642, 1584, 1529, 1505, 1403, 1309 cm^{-1} .

HRFABMS: Found 553.28890; calculated for $\text{C}_{27}\text{H}_{37}\text{N}_8\text{O}_5$ 553.28869.

(v) 1-Cyclopentyl-N-[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]-4-({[4-(formylamino)-1-methyl-1H-pyrrol-2-yl]-carbonyl}amino)-1H-pyrrole-2-carboxamide

1-Cyclopentyl-N-[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]-4-({[(1-methyl-4-nitro-1H-pyrrol-2-yl)carbonyl]amino}-1H-pyrrole-2-carboxamide (100 mg, 0.181 mmol; see step (iv) above) was dissolved in ethanol (20 mL) at 0°C under N_2 with stirring, to which Pd/C-10% (92 mg) was added. The reaction mixture was hydrogenated at room temperature and atmospheric pressure for 3 h. The catalyst was removed over Kieselguhr and ethyl formate (25 mL) was added to the ethanolic solution. The reaction mixture was heated under reflux for 24 h. The solvent was removed under reduced pressure to give the crude product, which was purified by HPLC. Fractions containing the required material were collected and freeze dried to give the product as pale yellow solid (21 mg, 18% yield) with no distinct melting point.

NMR [DMSO- d_6]: 10.06(1H, s); 9.95(1H, s); 9.91(1H, s); 9.21(1H, br, TFA); 8.14(1H, t, $J=6.0$ Hz); 8.13(1H, d, $J=1.7$ Hz); 7.36(1H, d, $J=1.7$ Hz); 7.19(1H, d, $J=1.7$ Hz); 7.17(1H, d, $J=1.7$ Hz); 6.97(1H, d, $J=1.7$ Hz); 6.95(1H, d, $J=1.7$ Hz); 6.92(1H, d, $J=1.7$ Hz); 5.52(1H, quintet, $J=7.5$ Hz);

3.84(3H, s); 3.81(3H, s); 3.24(2H, m); 3.06(2H, m); 2.79(6H, d, $J=3.3\text{Hz}$); 2.08(2H, m); 1.83(4H, m); 1.64(4H, m).

IR [KBr]: 1674, 1647, 1582, 1536, 1463, 1440, 1406, 1201, 1131 cm^{-1} .

HRFABMS: Found 551.30950; calculated for $\text{C}_{28}\text{H}_{39}\text{N}_8\text{O}_4$ 551.30943.

5

Example 28

N^2,N^7 -Bis[5-({[4-({[3-(dimethylamino)propyl]amino}carbonyl)-5-isopropyl-1,3-thiazol-2-yl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-9,10-dihydro-2,7-phenanthrenedicarboxamide

10 N -[3-(Dimethylamino)propyl]-5-isopropyl-2-{{[(1-methyl-4-nitro-1*H*-pyrrol-2-yl)carbonyl]amino}-1,3-thiazole-4-carboxamide (170 mg, 0.402 mmol; see Example 3, step (iv) above) was suspended in methanol (25 mL) at 0°C under N_2 with stirring, to which Pd/C-10% (104 mg) was added. The reaction mixture was hydrogenated for 6 h at room temperature and
15 atmospheric pressure. The catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure. The amine so formed was dissolved in DMF (2 mL, dry), to which 9,10-dihydro-2,7-phenanthrene-dicarboxylic acid (54 mg, 0.201 mmol; see *Tetrahedron* 56, 5225-5239 (2000)), HBTU (610 mg, 1.608 mmol) and NMM (200 μL , dry) were added
20 at room temperature with stirring. The reaction mixture was stirred overnight at room temperature. The reaction mixture was diluted with ethyl acetate containing 5% methanol and $\text{NaHCO}_3(\text{sat})$. The organic layer was dried over MgSO_4 , filtered and the solvent removed under reduced pressure to give the crude product, which was purified by HPLC. Fractions
25 containing the required material were collected and freeze dried to give pale yellow solid with distinct melting point (46.3 mg, 19% yield).

NMR [$\text{DMSO}-d_6$]: 12.11(2H, s); 10.46(2H, s); 9.26(2H, br, TFA); 8.07-7.91(6H, m); 7.53(2H, s); 7.46(2H, s); 4.20(2H, quintet, $J=6.9\text{Hz}$); 3.92(6H, s); 3.33(4H, m); 3.07(4H, m); 2.97(2H, s); 2.96(2H, s); 2.80(12H, d, $J=4.1\text{Hz}$); 1.87(4H, quintet, $J=7.7\text{Hz}$); 1.27(12H, d, $J=6.9\text{Hz}$).

30

IR [KBr]: 1660, 1548, 1468, 1284, 1199, 1132, 832, 800, 721 cm⁻¹.

LRESMS: Found M+H=1017.4; [M+2]/2=509.5; calculated for C₅₂H₆₆N₁₂O₆S₂ M+H=1017.4; [M+2H]/2=509.3.

5 Example 29

4-(Formylamino)-N-[1-isopentyl-5-({[1-methyl-5-({[3-(4-methyl-1-piperazinyl)propyl]amino}carbonyl)-1H-pyrrol-3-yl]amino}carbonyl)-1H-pyrrol-3-yl]-1-methyl-1H-pyrrole-2-carboxamide

- 10 (i) 1-Isopentyl-N-[1-methyl-5-({[3-(4-methyl-1-piperazinyl)propyl]amino}carbonyl)-1H-pyrrol-3-yl]-4-nitro-1H-pyrrole-2-carboxamide
1-Methyl-N-[3-(4-methyl-1-piperazinyl)propyl]-4-nitro-1H-pyrrole-2-carboxamide (209 mg, 0.676 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) was dissolved in methanol (25 mL) to which Pd/C-10% (182 mg) was
15 added at 0°C under N₂ with stirring. The reaction mixture was hydrogenated for 3 h at room temperature and atmospheric pressure. The catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure to give the amine, which was used without further purification. 1-Isopentyl-4-nitro-1H-pyrrole-2-carboxylic acid (153 mg,
20 0.676 mmol; prepared according to the procedure described in *Soviet Journal of Bioorganic Chemistry (English Translation)* 4, 780-790 (1978)) was dissolved in thionyl chloride 4 mL and heated under reflux for 3 h. Excess thionyl chloride was removed under pressure at 50°C and the acid chloride was dissolved in DCM (5 mL). The amine was dissolved in DCM
25 (10 mL) to which NMM (200 µL) was added followed by the acid chloride at room temperature with stirring. The stirring was continued at room temperature overnight. KOH 10% (5 mL) was added and the mixture was extracted. The organic layer was collected, dried over MgSO₄, filtered and the solvent was removed under reduced pressure to give the crude product,
30 which was purified by column chromatography using silica gel and 1/2/0.1

methanol/ethyl acetate/TEA. The product was obtained as glassy yellow material ($R_F=0.45$), (278 mg, 84% yield) with no distinct melting point.

NMR [DMSO- d_6]: 10.21(1H, s); 8.23(1H, d, $J=1.8$ Hz); 8.05(1H, t, $J=5.4$ Hz); 7.54(1H, d, $J=1.8$ Hz); 7.18(1H, d, $J=1.8$ Hz); 6.82(1H, d, $J=1.8$ Hz); 4.43(2H, t, $J=7.4$ Hz); 3.81(3H, s); 3.21-3.16(2H, q, $J=6.6$ Hz); 2.32-2.28(10H, m); 2.15(3H, s); 1.66-1.59(4H, quintet, $J=6.7$ Hz); 1.55-1.47(1H, m); 0.90(6H, d, $J=6.5$ Hz).

IR [KBr]: 2951, 2805, 1642, 1575, 1532, 1506, 1437, 1312 cm^{-1} .

HRFABMS: Found 488.29707; calculated for $\text{C}_{24}\text{H}_{38}\text{N}_7\text{O}_4$ 488.29853.

(ii) *N*-[1-Isopentyl-5-({[1-methyl-5-({[3-(4-methyl-1-piperazinyl)propyl]-amino}carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1-methyl-4-nitro-1*H*-pyrrole-2-carboxamide

1-Isopentyl-*N*-[1-methyl-5-({[3-(4-methyl-1-piperazinyl)propyl]amino}-carbonyl)-1*H*-pyrrol-3-yl]-4-nitro-1*H*-pyrrole-2-carboxamide (140 mg, 0.287 mmol; see step (i) above) was dissolved in methanol (25 mL) to which Pd/C-10% (100 mg) was added at 0°C under N_2 with stirring. The reaction mixture was hydrogenated at room temperature and atmospheric pressure for 3 h. The catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure. 1-Methyl-4-nitro-1*H*-pyrrole-2-carboxylic acid (49 mg, 0.287 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) was dissolved in thionyl chloride (4 mL) and the reaction mixture was heated under reflux for 3 h. Excess thionyl chloride was removed under reduced pressure to give the acid chloride, which was dissolved in DCM (5 mL). The amine was dissolved in DCM (10 mL) to which NMM (0.1 mL) was added at room temperature with stirring followed by the acid chloride solution. The stirring was continued overnight at room temperature. KOH (5 mL, 10%) was added and the reaction mixture was extracted, and the organic layer collected, dried over MgSO_4 , filtered and the solvent removed under reduced pressure. Column chromatography

(silica gel, 1/2/0.1 methanol/ethyl acetate/TEA) was used to purify the required material, which was obtained as yellow glassy material (131 gm, 75% yield), $R_f=0.15$ with no distinct melting point.

NMR [DMSO- d_6]: 10.27(1H, s); 9.90(1H, s); 8.18(1H, d, $J=1.8\text{Hz}$);
5 8.01(1H, t, $J=5.6\text{Hz}$); 7.59(1H, d, $J=1.8\text{Hz}$); 7.31(1H, d, $J=1.8\text{Hz}$); 7.16(1H, d, $J=1.8\text{Hz}$); 6.98(1H, $J=1.8\text{Hz}$); 6.84(1H, d, $J=1.8\text{Hz}$); 4.33(1H, t, $J=6.9\text{Hz}$); 3.96(3H, s); 3.79(3H, s); 3.17(2H, q, $J=6.5\text{Hz}$); 2.32-2.29(10mH, m); 2.16(3H, s); 1.66-1.46(5H, m); 0.89(6H, d, $J=6.4\text{Hz}$).

IR [KBr]: 2949, 2803, 1650, 1588, 1531, 1506, 1399, 1309 cm^{-1} .

10 HRFABMS: Found 610.34797; calculated for $\text{C}_{30}\text{H}_{44}\text{N}_9\text{O}_5$ 610.34654.

(iii) 4-(Formylamino)-N-[1-isopentyl-5-({[1-methyl-5-({[3-(4-methyl-1-piperazinyl)propyl]amino}carbonyl)-1H-pyrrol-3-yl]amino}carbonyl)-1H-pyrrol-3-yl]-1-methyl-1H-pyrrole-2-carboxamide

15 *N*-[1-Isopentyl-5-({[1-methyl-5-({[3-(4-methyl-1-piperazinyl)propyl]-amino}carbonyl)-1H-pyrrol-3-yl]amino}carbonyl)-1H-pyrrol-3-yl]-1-methyl-4-nitro-1H-pyrrole-2-carboxamide (120 mg, 0.197 mmol; see step (ii) above) was dissolved in ethanol (20 mL) to which Pd/C-10% (77 mg) was added at 0°C under N_2 with stirring. The reaction mixture was
20 hydrogenated for 3 h at room temperature and atmospheric pressure. The catalyst was removed over Kieselguhr and ethyl formate (20 mL) was added to the ethanolic solution. The reaction mixture was heated under reflux for 48 h. Ethanol and excess ethyl formate were removed under reduced pressure and the crude product was purified by HPLC. The product was
25 obtained as yellow solid with no distinct melting point (75 mg, 53% yield) after freeze-drying.

NMR [DMSO- d_6]: 10.03(1H, s); 9.89(1H, s); 9.87(1H, s); 8.13(1H, d, $J=1.6\text{Hz}$); 8.09(1H, t, $J=5.6\text{Hz}$); 7.26(1H, d, $J=1.6\text{Hz}$); 7.18(1H, d, $J=1.6\text{Hz}$); 7.14(1H, d, $J=1.6\text{Hz}$); 7.01(1H, d, $J=1.6\text{Hz}$); 6.92(1H, d, $J=1.6\text{Hz}$); 6.91(1H, d, $J=1.6\text{Hz}$); 4.31(2H, t, $J=6.9\text{Hz}$); 3.83(3H, s);
30

3.81(3H, s); 3.23(2H, q, J=6.5Hz); 3.10-2.72(8H, br & 3H, s); 1.77(2H, s, br); 1.58-1.46(3H, m); 0.90(6H, d, J=6.4Hz).

IR [KBr]: 1675, 1584, 1535, 1403, 1199, 1132 cm⁻¹.

HRFABMS: Found 608.36634; calculated for C₃₁H₄₆N₉O₄ 608.36728.

5

Example 30

4-(Acetylamino)-N-[1-isopentyl-5-({[1-methyl-5-({[3-(4-morpholinyl)-propyl]amino}carbonyl)-1H-pyrrol-3-yl]amino}carbonyl)-1H-pyrrol-3-yl]-1-methyl-1H-pyrrole-2-carboxamide

10

(i) 1-Isopentyl-N-[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}carbonyl)-1H-pyrrol-3-yl]-4-nitro-1H-pyrrole-2-carboxamide

1-Methyl-N-[3-(4-morpholinyl)propyl]-4-nitro-1H-pyrrole-2-carboxamide (500 mg, 1.688 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) was dissolved in methanol (25 mL) at 0°C under N₂ with stirring to which was added Pd/C-10% (272 mg). The reaction mixture was hydrogenated for 4 h. The catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure. The amine so formed was dissolved in DCM (10 mL). 1-Isopentyl-4-nitro-1H-pyrrole-2-carboxylic acid (445 mg, 1.966 mmol; prepared according to the procedure described in *Soviet Journal of Bioorganic Chemistry (English Translation)* 4, 780-790 (1978)) was dissolved in thionyl chloride (5 mL) and the reaction mixture was heated under reflux for 4 h. Excess thionyl chloride was removed under reduced pressure and the acid chloride was dissolved in DCM (10 mL). NMM (0.2 mL) was added to the amine solution followed by the acid chloride solution at room temperature with stirring. The stirring was continued at room temperature overnight. DCM was removed under reduced pressure and the crude product was purified by column chromatography using silica gel and 1/2/0.1 methanol/ethyl acetate/TEA. Fractions containing the pure material (R_F=0.45) were collected and the

solvents removed under reduced pressure to give yellow glassy material, which was dissolved in small amount of ethyl acetate and precipitated with *n*-hexane to give the required product as yellow powder (670 mg, 79% yield), m.p.155-158°C.

5 NMR [CDCl₃]: 7.64(1H, d, J=1.7Hz); 7.60(1H, s); 7.22(1H, unresolved triplet); 7.18(1H, d, J=1.7Hz); 7.07(1H, d, J=1.7Hz); 6.66(1H, d, J=1.6Hz); 4.43(2H, t, J=7.5Hz); 3.93(3H, s); 3.77(2H, t, J=4.6Hz); 3.49(2H, q, J=5.6Hz); 2.52(6H, m); 1.80-1.58(5H, m); 0.97(6H, d, J=6.5Hz).

IR [KBr]: 1647, 1589, 1513, 1399, 1309, 1252, 1114 cm⁻¹.

10 HRFABMS: Found 475.26789; calculated for C₂₃H₃₅N₆O₅ 475.26689.

(ii) *N*-[1-Isopentyl-5-({[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}-carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1-methyl-4-nitro-1*H*-pyrrole-2-carboxamide

15 1-Isopentyl-*N*-[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]-4-nitro-1*H*-pyrrole-2-carboxamide (136 mg, 0.286 mmol; see step (i) above) was dissolved in methanol (25 mL) at 0°C under N₂ with stirring to which was added Pd/C-10% (98 mg). The reaction mixture was hydrogenated at room temperature and atmospheric pressure for 2 h. The
20 catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure. The amine so formed was dissolved in DCM (5 mL). 1-Methyl-4-nitro-1*H*-pyrrole-2-carboxylic acid (48 mg, 0.282 mmol; see *Tetrahedron* 56, 5225-5239 (2000)) was dissolved in thionyl chloride (3 mL) and heated until reflux for 2 h. Excess thionyl chloride was
25 removed under reduced pressure and the acid chloride so formed was dissolved in DCM (5 mL) and then added dropwise to the amine solution with stirring at room temperature. The stirring was continued overnight. The solvent was removed and the crude product was purified by column chromatography (1/2/0.1 methanol/ethyl acetate/ TEA, R_F=0.7). Fractions
30 containing the required material were collected and the solvents were

removed under reduced pressure. The yellow glassy material was dissolved in DCM (25 mL) and extracted with KOH (10%, 5 mL). The organic layer was collected, dried over MgSO₄, filtered and the solvent removed under reduced pressure to give yellow glassy material (115 mg, 67% yield) with no distinct melting point.

NMR [Acetone-d₆]: 9.52(1H, s); 9.26(1H, s); 7.95(1H, d, J=1.7Hz); 7.55(1H, t, unresolved); 7.38(1H, d, J=1.9Hz); 7.33(1H, d, J=1.9Hz); 7.20(1H, d, J=1.9Hz); 6.91(1H, d, J=1.9Hz); 6.84(1H, d, J=1.9Hz); 4.44(2H, t, J=7.2Hz); 4.08(3H, s); 3.89(3H, s); 3.64(4H, m); 3.37(2H, t, J=6.7Hz); 2.42(6H, m); 1.77-1.56(5H, m); 0.95(6H, d, J=6.5Hz).

IR [KBr]: 1640, 1588, 1524, 1464, 1399, 1310, 1252, 1114 cm⁻¹.

HRFABMS: 597.31332; calculated for C₂₉H₄₁N₈O₆ 597.31491.

(iii) 4-(Acetylamino)-N-[1-isopentyl-5-({[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}carbonyl)-1H-pyrrol-3-yl]amino}carbonyl)-1H-pyrrol-3-yl]-1-methyl-1H-pyrrole-2-carboxamide

N-[1-Isopentyl-5-({[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}carbonyl)-1H-pyrrol-3-yl]amino}carbonyl)-1H-pyrrol-3-yl]-1-methyl-4-nitro-1H-pyrrole-2-carboxamide (115 mg, 0.193 mmol; see step (ii) above) was dissolved in ethanol (20 mL) to which was added Pd/C-10% (104 mg) at 0°C under N₂ with stirring. The reaction mixture was hydrogenated for 3 h at room temperature and atmospheric pressure. The catalyst was removed over Kieselguhr and the ethanolic solution was divided into two equal volumes (*i.e.* 10 mL). Ethanol was removed from one of the fractions under reduced pressure and DCM (2 mL) was added to the amine with stirring. NMM (0.1 mL) was added followed by acetyl chloride (0.01 mL) at room temperature with stirring. The stirring was continued overnight. All volatile materials were removed under reduced pressure and the crude product was purified by HPLC. Fractions containing the required material

were collected and freeze-dried to give white solid material (25.7 mg, 35% yield) as TFA salt with no distinct melting point.

NMR [DMSO- d_6]: 9.87(2H, s); 9.79(1H, s); 9.55(1H, br, TFA); 8.16(1H, t, unresolved); 7.26(1H, d, $J=1.8\text{Hz}$); 7.15(1H, d, $J=1.8\text{Hz}$); 7.13(1H, d, $J=1.8\text{Hz}$); 7.01(1H, d, $J=1.8\text{Hz}$); 6.94(1H, d, $J=1.8\text{Hz}$); 6.87(1H, d, $J=1.8\text{Hz}$); 4.31(2H, t, $J=6.9\text{Hz}$, isopentyl); 4.00(2H, d, $J=12.4\text{Hz}$, morpho); 3.83(3H, s); 3.81(3H, s); 3.64(2H, t, $J=12.0\text{Hz}$, morpho); 3.45(2H, d, $J=12.0\text{Hz}$, morpho); 3.24(2H, q, $J=6.2\text{Hz}$); 3.12-3.06(2H, m, side chain & 2H morpho); 1.97(3H, s); 1.87(2H, m, isopentyl); 1.58-1.48(2H, m, side chain & 1H isopentyl); 0.90(6H, d, $J=6.4\text{Hz}$, isopentyl).

IR [KBr]: 2925, 2858, 1655, 1583, 1526, 1465, 1400, 1260, 1200, 1131 cm^{-1} .

HRFABMS: Found 609.34957; calculated for $\text{C}_{31}\text{H}_{45}\text{N}_8\text{O}_5$ 609.35129.

15 Example 31

4-(Formylamino)-*N*-[1-isopentyl-5-({[1-methyl-5-({[3-(4-morpholinyl)-propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1-methyl-1*H*-pyrrole-2-carboxamide

To the second half of the amine solution (10 mL) (see Example 30, step (iii) above), ethyl formate was added (10 mL) and the reaction mixture was heated under reflux for 48 h. Volatile solvents were removed under reduced pressure and the crude product was purified by HPLC. Fractions containing the required material were collected and freeze-dried to give white solid material (22.7 mg, 32% yield) as TFA salt with no distinct melting point.

25 NMR [DMSO- d_6]: 10.03(1H, s); 9.89(1H, s); 9.87(1H, s); 9.56(1H, br, TFA); 8.16(1H, t, unresolved); 8.12(1H, d, $J=1.7\text{Hz}$); 7.26(1H, d, $J=1.7\text{Hz}$); 7.18(1H, d, $J=1.7\text{Hz}$); 7.15(1H, d, $J=1.7\text{Hz}$); 7.02(1H, d, $J=1.7\text{Hz}$); 6.93(1H, d, $J=1.7\text{Hz}$); 6.92(1H, d, $J=1.7\text{Hz}$); 4.31(2H, t, $J=7.1\text{Hz}$); 4.00(2H, d, $J=11.8\text{Hz}$); 3.84(3H, s); 3.81(3H, s); 3.64(2H, t, $J=11.9\text{Hz}$); 3.46(2H, d,

J=12.8Hz); 3.26(2H, q, J=6.1Hz); 3.12-3.06(4H, m); 1.87(2H, m); 1.58-1.46(3H, m); 0.90(6H, d, J=6.3Hz).

IR [KBr]: 1682, 1640, 1584, 1529, 1403, 1263, 1202, 1132, 803 cm⁻¹.

HRFABMS: Found 595.33717; calculated for C₃₀H₄₃N₈O₅ 595.33564.

5

Example 32

N-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1H-pyrrol-3-yl]-4-[(3-methoxybenzoyl)amino]-1-methyl-1H-pyrrole-2-carboxamide

10 N-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1H-pyrrol-3-yl]-1-methyl-4-nitro-1H-pyrrole-2-carboxamide (260 mg, 0.469 mmol; see Example 10, step (ii) above) was dissolved in methanol (25 mL) to which Pd/C-10% (132 mg) was added at 0°C under N₂ with stirring. The reaction mixture was
15 hydrogenated for 3 h. The catalyst was removed over Kieselguhr and the solvent was removed under reduced pressure to give the amine, which was dissolved in DCM (5 mL). This solution was divided to two each (2.5 mL). *m*-Anisoyl chloride (50 mg, 0.293 mmol) was added dropwise at room temperature with stirring. The stirring was continued overnight. The
20 solvent was removed under reduced pressure and the residue was dissolved in acetonitrile containing (0.1%TFA) and purified by HPLC. Fractions containing the required material were collected and freeze-dried to give the product as light pink solid with no distinct melting point (72.3 mg, 40% yield).

25 NMR [DMSO-d₆]: 10.29(1H, s); 9.96(1H, s); 9.89(1H, s); 9.27(1H, br, TFA); 8.15(1H, t, J=5.6Hz); 7.53-7.37(3H, m); 7.32(1H, d, J=1.6Hz); 7.29(1H, d, J=1.6Hz); 7.16(1H, d, J=1.6Hz); 7.14-7.10(2H, m); 7.02(1H, d, J=1.6Hz); 6.93(1H, d, J=1.6Hz); 4.32(1H, t, J=6.9Hz); 3.88(3H, s); 3.83(3H, s); 3.82(3H, s); 3.25(2H, q, J=6.2Hz); 3.07(2H, m); 2.79(6H, d,

J=4.5Hz); 1.84(2H, quintet, J=7.3Hz); 1.59-1.47(3H, m); 0.90(6H, d, J=6.3Hz).

IR [KBr]: 1644, 1583, 1533, 1464, 1436, 1402, 1260, 1200, 801, 778 cm⁻¹.

HRFABMS: Found 659.36784; calculated for C₃₅H₄₇N₈O₅ 659.36694.

5

Example 33

N-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1H-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1H-pyrrol-3-yl]-4-{{[(4-methoxy-phenyl)acetyl]amino}-1-methyl-1H-pyrrole-2-carboxamide

10 The second half of the amine (see Example 32 above) was used in this experiment. To the amine (2.5 mL) in DCM was added (4-methoxy-phenyl)acetyl chloride (50 mg, 0.270 mmol) at room temperature, dropwise with stirring. The stirring was continued overnight. The solvent was removed under reduced pressure and the crude product so formed was
15 purified by HPLC. Fractions containing the required material were collected and freeze-dried to give white solid material (51 mg, 28% yield) with no distinct melting point.

NMR [Acetone-d₆]: 12.15(1H, br); 9.18(1H, s); 9.11(1H, s); 9.05(1H, s); 7.67(1H, t, unresolved); 7.29-7.24(4H, m); 7.13(1H, d, J=1.7Hz); 6.89-
20 6.85(4H, m); 6.78(1H, d, J=1.7Hz); 4.41(2H, t, J=6.5Hz); 3.90(6H, s); 3.77(3H, s); 3.58(2H, s); 3.45(2H, q, J=6.1Hz); 3.26(2H, t, J=6.1Hz); 2.93(6H, s); 1.69-1.56(3H, m); 0.95(6H, d, J=6.6Hz).

IR [KBr]: 1650, 1588, 1515, 1465, 1402, 1251, 1203, 1133, 826, 779 cm⁻¹.

HRFABMS: Found 673.38492; calculated for C₃₆H₄₉N₈O₅ 673.38259.

25

Example 34

Compounds of the examples were found to bind to the minor groove of DNA, as determined by the capillary electrophoresis and/or DNA footprinting methods described hereinbefore.

Example 35

Experiments that probe the binding of compounds to DNA by using n.m.r. methods are well known to those skilled in the art (see, for example: (a) J. Bunkenborg, C. Behrens, J. P. Jacobsen "*NMR characterization of the DNA binding properties of a novel Hoechst 33258 analogue peptide building block*" *Bioconjugate Chemistry* 13(5), 927-936 (2002); (b) G. A Morris, K. T. Douglas "*Binding of a porphyrin conjugate of Hoechst 33258 to DNA. II. NMR spectroscopic studies detect multiple binding modes to a 12-mer nonself-complementary duplex DNA*" *Nucleosides Nucleotides & Nucleic Acids* 20(1-2), 145-156 (2001); and (c) X. G. Han, X. L. Gao "*Sequence specific recognition of ligand-DNA complexes studied by NMR*" *Current Medicinal Chemistry* 8(5) 551-581 (2001)).

Compounds of the examples were found to bind to the minor groove of DNA, as determined by experiments in which the compounds were mixed under standard conditions with DNA duplexes (e.g. d(CGACTAGTCG)₂) and binding was confirmed by changes observed in the n.m.r. spectra of the compound and the DNA duplex, as well as by nOe measurements on the complex formed between the compound and the DNA duplex. Narrow lines and strong cross-peaks were observed for the complex formed between the compound of Example 3 and d(CGACTAGTCG)₂, which is consistent with the compound of Example 3 binding with high affinity.

Example 36

Compounds of the examples were found to bind to DNA, as determined by melting temperature (T_m) measurements. T_m measurements are well known in the art as a method for investigating the stability of a DNA duplex (see V. A Bloomfield, D. M. Crothers and I. Tinoco "*Nucleic Acids: Structures, Properties and Functions*" University Science Books, Sausalito, California (2000), and in particular pages 176-180 and 561-564 of that

book). If a ligand stabilises the duplex, an increase in T_m is recorded. Compounds of the examples demonstrated large increases in T_m for DNA duplexes. For example, when the DNA used was a duplex of AAATTATATTAT, the compounds of Examples 2, 10 and 26 gave an
 5 increase in melting point of greater than 10°C.

Example 37

Compounds of the examples were found to inhibit the growth of microorganisms, for example as indicated in Tables A to G below.

10

Table A

Example No.	MIC (μM) Organism: S. aureus	MIC (μM) Organism: E. coli
7	4.8	>152.4
17	19.4	>77.7
28	80	>80.3
3	4.7	>152.3
Antibiotic control Amoxicillin	0.49	2.0

Table B

Example No.	MIC (μM) Organism: S.faecalis	MIC (μM) Organism: P.vulgaris
7	9.5	152.4
17	77.7	77.7
28	10.0	>80.3
3	9.5	152.2
Antibiotic control Amoxicillin	0.49	8.1

Table C

Example No.	MIC (μM) Organism: MRSA
7	19.1
17	>77.7
28	10.0
3	38.1
Antibiotic control Amoxicillin	16.1

Table D

Example No.	MIC (μM) Organism: <i>Aspergillus niger</i>	MIC (μM) Organism: <i>Candida albicans</i>
7	38.1	38.1
17	>77.7	>77.7
28	>80.3	>80.3
3	76.1	76.1
Anti fungal control Fluconazole	>326.6	81.6

Table E

Example No.	MIC (μM) Organism: <i>Klebs.</i> <i>aerogenes</i>	MIC (μM) Organism: <i>Ent. cloacae</i>
7	38.1	9.5
17	38.9	>77.7
28	>80.3	>80.3
3	76.2	76.2
Control Amoxicillin	32.3	4.0

Table F

Example No.	MIC (μM) Organism: <i>Mycobacterium fortuitum</i>
7	19.0
17	>77.7
28	40.1
3	19.0
Control Streptomycin	21.5

Table G

Example No.	MIC (μM) Organism: <i>Aspergillus nidulans</i>
7	9.5
17	77.7
28	20.1
3	38.1
Control Fluconazole	40.8

5 Abbreviations

br	=	broad (in relation to NMR)
CE	=	capillary electrophoresis
d	=	doublet (in relation to NMR)
10 DCM	=	dichloromethane
DMF	=	<i>N,N</i> -dimethylformamide

	DMSO	=	dimethylsulfoxide
	eq.	=	equivalents
	h	=	hour(s)
	HBTU	=	<i>O</i> -benzotriazol-1-yl- <i>N,N,N',N'</i> -tetramethyluronium
5			hexafluorophosphate
	HCl	=	hydrochloric acid
	HOBT	=	1-hydroxybenzotriazole
	HPLC	=	high performance liquid chromatography
	HREIMS	=	high resolution electron ionisation mass spectrometry
10	HRFABMS	=	high resolution fast atom bombardment mass spectrometry
	IR	=	infra red (in relation to spectroscopy)
	LRESMS	=	low resolution electrospray mass spectrometry
	m	=	multiplet (in relation to NMR)
15	Me	=	methyl
	min.	=	minute(s)
	m.p.	=	melting point
	MS	=	mass spectroscopy
	ν_{\max}	=	wave number (in relation to infra red spectroscopy)
20	NMM	=	<i>N</i> -methyldmorpholine
	Pd/C	=	palladium on carbon
	q	=	quartet (in relation to NMR)
	s	=	singlet (in relation to NMR)
	t	=	triplet (in relation to NMR)
25	TEA	=	triethylamine
	THF	=	tetrahydrofuran
	TFA	=	trifluoroacetic acid

Prefixes *n*-, *s*-, *i*-, *t*- and *tert*- have their usual meanings: normal, secondary,
30 *iso*, and tertiary.


Claims

1. An oligopeptide compound comprising:
 - (a) at least one nitrogen-containing basic group attached to at least
5 one end of the oligopeptide; and
 - (b) two or more heterocyclic monomers, at least one of which is substituted in the heterocyclic part by a branched, cyclic or part cyclic C₃₋₅ alkyl group,
or a pharmaceutically acceptable salt or solvate thereof;
10 which compound, salt or solvate binds to the minor groove of DNA.
2. An oligopeptide compound comprising:
 - (a) at least one nitrogen-containing basic group attached to at least
one end of the oligopeptide; and
 - 15 (b) two or more heterocyclic monomers, at least one of which is substituted on a ring C-atom in the heterocyclic part by a branched, cyclic or part cyclic C₃₋₅ alkyl group,
or a pharmaceutically acceptable salt or solvate thereof;
which compound, salt or solvate binds to the minor groove of DNA.
20
3. A compound as claimed in Claim 1 or Claim 2, wherein the nitrogen-containing basic group, in its neutral state, has a pK_a in water of at least 4.
4. A compound as claimed in Claim 3, wherein the nitrogen-containing
25 basic group is an amidino, guanidino or amino group, any of which may be cyclic or acyclic.
5. A compound as claimed in any one of the preceding claims, wherein
each heterocyclic group is a 4- to 12-membered heterocyclic group
30 containing one or more heteroatoms selected from N, O and S.

6. A compound as claimed in Claim 5, wherein each heterocyclic group is independently aromatic or part-aromatic.
7. A compound as claimed in Claim 6, wherein each heterocyclic group that is present is independently selected from the group consisting of pyrrolyl, imidazolyl, thiazolyl, oxazolyl, benzoxazolyl, furanyl, thienyl, pyridyl and coumarinyl.
8. A compound as claimed in Claim 7, wherein at least one heterocyclic group that is present is a thiazolyl group.
9. A compound as claimed in Claim 8, wherein the thiazolyl group is a 1,3-thiazolyl group that is substituted in the 5-position by, as appropriate, the branched, cyclic or part cyclic C₃₋₅ alkyl group.
10. A compound as claimed in any one of the preceding claims, wherein the branched, cyclic or part cyclic C₃₋₅ alkyl group is isopropyl.
11. A compound as claimed in any one of the preceding claims, which has a molecular weight of below 2000 g mol⁻¹.
12. A compound as claimed in any one of the preceding claims, which is bioavailable.
13. A compound as claimed in any one of the preceding claims, which has a high affinity for DNA sequences.
14. A compound as claimed in Claim 12 that binds to the minor groove of a DNA oligomer or polymer with a dissociation constant of less than 10⁻⁵ M.

15. A compound as claimed in Claim 1, wherein each essential branched, cyclic or part cyclic C₃₋₅ alkyl substituent on a heterocyclic monomer is other than cyclopropyl.

5 16. A compound of formula I,



wherein

R¹ represents Het¹, R^{1a}C(O)- or D-A-N(H)-[Q]_n-C(O)-E-C(O)-;

R^{1a} represents

10 H,

aryl (which latter group is optionally substituted by one or more substituents selected from OH, halo, cyano, nitro, N(R^{3a})R^{3b}, C₁₋₄ alkyl and C₁₋₄ alkoxy),

15 aromatic or part-aromatic C₁₃₋₁₄ tricyclic carbocyclyl (which latter group is optionally substituted by one or more substituents selected from OH, halo, cyano, nitro, N(R^{3a})R^{3b}, C₁₋₄ alkyl and C₁₋₄ alkoxy, and which latter group, if part-aromatic, is optionally substituted in the non-aromatic part by one or two oxo groups) or

20 C₁₋₁₂ alkyl (which latter group is optionally substituted and/or terminated by one or more substituents selected from halo and aryl (which latter group is optionally substituted by one or more substituents selected from OH, halo, cyano, nitro, N(R^{3a})R^{3b}, C₁₋₄ alkyl and C₁₋₄ alkoxy));

25 A represents, at each occurrence when used herein, C₂₋₆ alkylene or A¹-C(O)N(H)-A², wherein A² is attached to the group D;

A¹ represents C₁₋₄ alkylene;

A² represents C₂₋₅ alkylene;

127


D represents, at each occurrence when used herein, $-N(R^{2a})R^{2b}$, $-C(=NR^{2c})N(R^{2d})R^{2e}$ or $-N(R^{2f})C(=NR^{2g})N(H)R^{2h}$;

R^{2a} and R^{2b} independently represent H, C_{1-6} alkyl, Het² or R^{2a} and R^{2b} together represent $(CH_2)_{3-6}$, which alkylene group is optionally interrupted by NR^4 and/or is optionally substituted by one or more C_{1-4} alkyl groups;

R^4 represents H, C_{1-6} alkyl or Het³;

R^{2c} to R^{2h} independently represent H or C_{1-6} alkyl;

E represents $-E^1-Het^4-$, E^{2a} , $-(CH_2)_{0-3}N(H)C(O)-E^{2b}-C(O)N(H)(CH_2)_{0-3}-$ or a structural fragment of the formula



wherein E^3 represents $(CH_2)_{1-2}$, $CH=CH$, $CH=N$, $CH_2-N(R^a)$, $(CH_2)_{0-1}C(O)$, $(CH_2)_{0-1}O$ or $(CH_2)_{0-1}S$;

R^a represents H or C_{1-6} alkyl;

E^1 represents $(CH_2)_{0-2}$ or $CH=CH$;

E^{2a} and E^{2b} independently represent C_{2-4} alkenylene, C_{3-6} cycloalkylene, phenylene or naphthylene;


Het¹ to Het⁴ independently represent four- to twelve-membered heterocyclic groups containing one or more heteroatoms selected from N, O and S, which heterocyclic groups are optionally substituted by one or more substituents selected from =O, OH, halo, cyano, nitro, $N(R^{3a})R^{3b}$, C_{1-4} alkyl and C_{1-4} alkoxy;

R^{3a} and R^{3b} independently represent, at each occurrence when used herein, H or C_{1-4} alkyl, or R^{3a} represents $-C(O)R^5$;


R^5 represents H or C_{1-4} alkyl;

n represents, at each occurrence when used herein, 2, 3, 4 or 5;


each individual Q independently represents a structural fragment of formula Ia, Ib, Ic, Id, Ie or If




Ia




Ib




Ic



Id



Ie



If

wherein

R⁶ represents H or C₁₋₆ alkyl;

10 R⁷ represents C₁₋₁₂ alkyl;

R⁸, R⁹, R¹⁰ and R¹¹ independently represent H or C₁₋₁₂ alkyl;

G represents CH or N;

L represents O or S;

p, q and r independently represent 0, 1, 2 or 3; and

provided that the compound comprises at least one structural fragment of formula Ib, Ic, Id, Ie or If in which R⁶ or R⁷, R⁸, R⁹, R¹⁰ or R¹¹, respectively, represents branched, cyclic or part cyclic C₃₋₅ alkyl;

5 or a pharmaceutically acceptable derivative thereof.

17. A compound as claimed in Claim 16, wherein:

10 R^{1a} represents H or C₁₋₁₂ alkyl, which latter group is optionally substituted and/or terminated by one or more substituents selected from halo and aryl, which latter group is optionally substituted by one or more substituents selected from OH, halo, cyano, nitro, N(R^{3a})R^{3b}, C₁₋₄ alkyl and C₁₋₄ alkoxy; and

15 the compound comprises at least one structural fragment of formula Ib, Ic, Id, Ie or If in which R⁷, R⁸, R⁹, R¹⁰ or R¹¹, respectively, represents branched, cyclic or part cyclic C₃₋₅ alkyl.

18. A compound as claimed in Claim 16 or Claim 17, wherein aryl is phenyl or naphthyl.

20 19. A compound as claimed in any one of Claims 16 to 18, wherein alkyl and alkoxy groups are, where appropriate:


- (a) straight-chain;
- (b) branched-chain and/or cyclic; or
- (c) part cyclic/acyclic.

25

20. A compound as claimed in any one of Claims 16 to 19, wherein alkyl and alkoxy groups are, where appropriate:

- (a) saturated or unsaturated;
- (b) interrupted by one or more oxygen and/or sulfur atoms; and/or
- 30 (c) unless otherwise specified, substituted by one or more halo atoms.

21. A compound as claimed in any one of Claims 16 to 20, which is a compound of formula II,



wherein

- 5 R^1 represents Het^1 , $\text{R}^{1a}\text{C(O)-}$ or $\text{D-A-N(H)-Q}^3\text{-Q}^2\text{-Q}^1\text{-C(O)-E-C(O)-}$;
 Q^1 is absent or represents a structural fragment of formula Ia, Ib, Ic, Id, Ie or If;
 Q^2 represents a structural fragment of formula Ib, Ie or If;
 Q^3 represents a structural fragment of formula Ib, Id, Ie or If; and
 10 Het^1 , R^{1a} , D, A, E, R^{2a} , R^{2b} , A and the structural fragments of formulae Ia, Ib, Ic, Id, Ie and If are as defined in any one of Claims 16 to 20;

provided that:

- (a) at least one of Q^1 , Q^2 and Q^3 represents a structural fragment of
 15 formula Id, Ie or If; and
 (b) at least one of R^6 or R^7 , R^8 , R^9 , R^{10} and R^{11} (whichever is/are present) represents branched, cyclic or part cyclic C_{3-5} alkyl,

or a pharmaceutically acceptable derivative thereof.

20

22. A compound as claimed in any one of Claims 16 to 21, wherein the compound comprises:


- (a) at least one structural fragment of formula Ib in which G represents N and R^6 represents branched, cyclic or part cyclic C_{3-5} alkyl;
 25 (b) at least one structural fragment of formula Id in which p represents 0 and R^9 represents branched, cyclic or part cyclic C_{3-5} alkyl; and/or
 (c) at least one structural fragment of formula Ie in which q represents 0 and R^{10} represents branched, cyclic or part cyclic C_{3-5} alkyl.

23. A compound as claimed in any one of Claims 16 to 23, wherein each of the at least one branched, cyclic or part cyclic C₃₋₅ alkyl groups independently represents isopropyl, cyclopropylmethyl, isopentyl or cyclopentyl.

5

24. A compound as claimed in any one of Claims 16 to 21, wherein the compound comprises at least one structural fragment of formula Ib, Ic, Id, Ie or If in which R⁷, R⁸, R⁹, R¹⁰ or R¹¹, respectively, represents isopropyl.

10 25. A compound as claimed in any one of Claims 16 to 24, which compound comprises at least one structural fragment of the formula



15 26. A compound as claimed in Claim 16, which compound is selected from the following:

(i) *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopropyl-1-*H*-pyrrol-3-yl]-4-[(3,3-dimethylbutanoyl)amino]-1-methyl-1*H*-pyrrole-2-carboxamide;

20 (ii) *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopropyl-1-*H*-pyrrol-3-yl]-4-(formylamino)-1-methyl-1*H*-pyrrole-2-carboxamide;

(iii) *N*-[3-(Dimethylamino)propyl]-2-({[4-({[4-(formylamino)-1-methyl-1*H*-pyrrol-2-yl]carbonyl}amino)-1-methyl-1*H*-pyrrol-2-yl]carbonyl}-amino)-5-isopropyl-1,3-thiazole-4-carboxamide;

- (iv) *N*-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopropyl-1*H*-pyrrol-3-yl]-4-({[4-(formylamino)-1-isopropyl-1*H*-pyrrol-2-yl]carbonyl}-amino)-1-isopropyl-1*H*-pyrrole-2-carboxamide
- (v) *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]-4-(formylamino)-1-isopentyl-1*H*-pyrrole-2-carboxamide;
- (vi) *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-isopropyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-4-(formylamino)-1-isopropyl-1*H*-pyrrole-2-carboxamide;
- (vii) *N*-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-2-({[4-(formylamino)-1-methyl-1*H*-pyrrol-2-yl]carbonyl}-amino)-5-isopropyl-1,3-thiazole-4-carboxamide;
- (viii) 4-({[4-(Formylamino)-1-methyl-1*H*-pyrrol-2-yl]carbonyl}amino)-1-iso-propyl-*N*-[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1*H*-pyrrole-2-carboxamide;
- (ix) 4-(Formylamino)-*N*-[1-isopropyl-5-({[1-methyl-5-({[3-(1-pyrrolidinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1-methyl-1*H*-pyrrole-2-carboxamide;
- (x) *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]-4-(formylamino)-1-methyl-1*H*-pyrrole-2-carboxamide;
- (xi) 2-(Acetylamino)-*N*-[5-({[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-5-isopropyl-1,3-thiazole-4-carboxamide;
- (xii) 2-(Acetylamino)-*N*-[5-({[4-({[3-(dimethylamino)propyl]amino}carbonyl)-5-isopropyl-1,3-thiazol-2-yl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-5-isopropyl-1,3-thiazole-4-carboxamide;
- (xiii) 2-(Acetylamino)-*N*-(5-({[3-({[3-(dimethylamino)propyl]amino}-3-oxo-propyl)amino]carbonyl}-1-methyl-1*H*-pyrrol-3-yl)-5-isopropyl-1,3-thiazole-4-carboxamide;

- (xiv) N^1, N^3 -Bis(2-{[5-({[4-({[3-(dimethylamino)propyl]amino}carbonyl)-5-isopropyl-1,3-thiazol-2-yl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-amino}-2-oxoethyl)isophthalamide;
- (xv) N -[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopropyl-1*H*-pyrrol-3-yl]-4-(acetylamino)-1-methyl-1*H*-pyrrole-2-carboxamide;
- (xvi) N -[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]-4-(acetylamino)-1-methyl-1*H*-pyrrole-2-carboxamide;
- (xvii) N^2, N^5 -Bis[5-({[4-({[3-(dimethylamino)propyl]amino}carbonyl)-5-isopropyl-1,3-thiazol-2-yl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-1*H*-indole-2,5-dicarboxamide;
- (xviii) N^2, N^5 -Bis[1-isopentyl-5-({[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1*H*-indole-2,5-dicarboxamide;
- (xix) N^2, N^5 -Bis[5-({[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]-1*H*-indole-2,5-dicarboxamide;
- (xx) N^2, N^5 -Bis[1-isopentyl-5-({[1-methyl-5-({[3-(4-methyl-1-piperazinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1*H*-indole-2,5-dicarboxamide;
- (xxi) 2-({[4-({[4-(Acetylamino)-1-methyl-1*H*-imidazol-2-yl]carbonyl}-amino)-1-methyl-1*H*-pyrrol-2-yl]carbonyl}amino)- N -[3-(dimethylamino)-propyl]-5-isopropyl-1,3-thiazole-4-carboxamide;
- (xxii) 4-(Acetylamino)- N -[1-isopentyl-5-({[1-methyl-5-({[3-(4-methyl-1-piperazinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1-methyl-1*H*-pyrrole-2-carboxamide;
- (xxiii) N -[1-Isopentyl-5-({[1-methyl-5-({[3-(4-methyl-1-piperazinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-4-[(3-methoxybenzoyl)amino]-1-methyl-1*H*-pyrrole-2-carboxamide;

- (xxiv) *N*-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-4-({[5-(formylamino)-2-methyl-3-thienyl]carbonyl}amino)-1-isopentyl-1*H*-pyrrole-2-carboxamide;
- (xxv) *N*-[5-({[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-5-isopropyl-2-[(3-methoxybenzoyl)amino]-1,3-thiazole-4-carboxamide;
- (xxvi) *N*-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-4-{{[5-{{[9,10-dioxo-9,10-dihydro-2-anthracenyl]carbonyl]-amino}-2-methyl-3-thienyl]carbonyl}amino}-1-isopentyl-1*H*-pyrrole-2-carboxamide;
- (xxvii) *N*-[1-(Cyclopropylmethyl)-5-({[5-({[3-(dimethylamino)propyl]-amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-4-(formylamino)-1-methyl-1*H*-pyrrole-2-carboxamide;
- (xxviii) 1-Cyclopentyl-*N*-[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-4-({[4-(formylamino)-1-methyl-1*H*-pyrrol-2-yl]-carbonyl}-amino)-1*H*-pyrrole-2-carboxamide;
- (xxix) *N*²,*N*⁷-Bis[5-({[4-({[3-(dimethylamino)propyl]amino}carbonyl)-5-isopropyl-1,3-thiazol-2-yl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-9,10-dihydro-2,7-phenanthrenedicarboxamide;
- (xxx) 4-(Formylamino)-*N*-[1-isopentyl-5-({[1-methyl-5-({[3-(4-methyl-1-piperazinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1-methyl-1*H*-pyrrole-2-carboxamide;
- (xxxi) 4-(Acetylamino)-*N*-[1-isopentyl-5-({[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1-methyl-1*H*-pyrrole-2-carboxamide;
- (xxxii) 4-(Formylamino)-*N*-[1-isopentyl-5-({[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1-methyl-1*H*-pyrrole-2-carboxamide;

(xxxiii) *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]-4-[(3-methoxybenzoyl)amino]-1-methyl-1*H*-pyrrole-2-carboxamide; and

(xxxiv) *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]-4-[(4-methoxyphenyl)acetyl] amino}-1-methyl-1*H*-pyrrole-2-carboxamide.

27. A compound as claimed in Claim 26 which is:

(a) *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopropyl-1*H*-pyrrol-3-yl]-4-(formylamino)-1-methyl-1*H*-pyrrole-2-carboxamide;

(b) *N*-[3-(Dimethylamino)propyl]-2-({[4-({[4-(formylamino)-1-methyl-1*H*-pyrrol-2-yl]carbonyl}amino)-1-methyl-1*H*-pyrrol-2-yl]carbonyl}-amino)-5-isopropyl-1,3-thiazole-4-carboxamide;

(c) *N*-[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-2-({[4-(formylamino)-1-methyl-1*H*-pyrrol-2-yl]carbonyl}-amino)-5-isopropyl-1,3-thiazole-4-carboxamide;

(d) *N*-[5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-isopentyl-1*H*-pyrrol-3-yl]-4-(formylamino)-1-methyl-1*H*-pyrrole-2-carboxamide;

(e) *N*²,*N*⁵-Bis[1-isopentyl-5-({[1-methyl-5-({[3-(4-morpholinyl)propyl]amino}carbonyl)-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-1*H*-indole-2,5-dicarboxamide;

(f) *N*-[1-(Cyclopropylmethyl)-5-({[5-({[3-(dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1*H*-pyrrol-3-yl]-4-(formylamino)-1-methyl-1*H*-pyrrole-2-carboxamide; or

(g) *N*²,*N*⁷-Bis[5-({[4-({[3-(dimethylamino)propyl]amino}carbonyl)-5-isopropyl-1,3-thiazol-2-yl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]-9,10-dihydro-2,7-phenanthrenedicarboxamide.

28. A compound as claimed in Claim 27 which is *N*-[3-(dimethylamino)-propyl]-2-({[4-({[4-(formylamino)-1-methyl-1*H*-pyrrol-2-yl]carbonyl}-amino)-1-methyl-1*H*-pyrrol-2-yl]carbonyl}amino)-5-isopropyl-1,3-thiazole-4-carboxamide.

5

29. A compound as claimed in any one of Claims 1 to 15, which binds to and/or has specificity for DNA sequences that contain at least one GC base pairing.

10 30. A compound as claimed in Claim 29, which is:

- (i) a compound of formula I, as defined in any one of Claims 16 to 20 or 22 to 25, provided that the compound comprises at least one structural fragment of formula Id, Ie or If; or
- (ii) a compound of formula II, as defined in any one of Claims 21 to 25.

15

31. A compound as claimed in any one of Claims 1 to 25 which has different binding affinities at different minor groove binding sites in double-stranded DNA molecules having more than one minor groove binding site.

20 32. A compound as claimed in Claim 31, wherein the different minor groove binding sites comprise solely AT base pairs.

33. A pharmaceutical formulation including a compound as defined in any one of Claims 1 to 32 in admixture with a pharmaceutically-acceptable
25 adjuvant, diluent or carrier.

34. A pharmaceutical formulation for use in the treatment of a disease that relies upon DNA replication for its propagation, comprising a compound as defined in any one of Claims 1 to 32 in admixture with a pharmaceutically-
30 acceptable adjuvant, diluent or carrier.

35. A compound as defined in any one of Claims 1 to 32 for use as a pharmaceutical.

36. A compound as defined in any one of Claims 1 to 32 for use in the
5 treatment of a disease that relies upon DNA replication for its propagation.

37. The use of a compound as defined in any of one Claims 1 to 32, as active ingredient for the manufacture of a medicament for use in the treatment of a disease that relies upon DNA replication for its propagation.

10

38. A method of treatment of a disease that relies upon DNA replication for its propagation, which method comprises administration of a therapeutically effective amount of a compound as defined in any of one Claims 1 to 32 to a person suffering from that disease.

15

39. A method of treating a viral, bacterial, fungal or other microbial (e.g. parasitic) infection, where the viral, bacterial, fungal or other microbial (e.g. parasitic) infective agent is resistant to one or more anti-viral, anti-bacterial, anti-fungal or other anti-microbial (e.g. anti-parasitic) agents, respectively,
20 that do not act by inhibiting DNA replication, which method comprises administration of a therapeutically effective amount of a compound as defined in any one of Claims 1 to 32 to a person having that infection.


40. A method of treatment of a disease that relies upon DNA replication
25 for its propagation, which method comprises administration, to a person suffering from that disease, of a therapeutically effective amount of a compound as defined in any one of Claims 1 to 32 in combination with one or more other agents that are known to be effective in treating that disease.

41. A combination product comprising components:
- (A) a formulation comprising a compound as defined in any one of Claims 1 to 32; and
 - (B) a formulation comprising one or more other chemical agents that are known to be effective in treating diseases that rely upon DNA replication for their propagation.
42. A combination product as claimed in Claim 41, wherein each of components (A) and (B) is formulated in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier.
43. A combination product as claimed in Claim 41 or Claim 42, wherein (A) and (B) are presented as separate components.
43. A combination product as claimed in Claim 41 or Claim 42, wherein (A) and (B) are presented as a single formulation.
45. A method of inhibiting DNA replication, which method comprises contacting the DNA with an inhibitory amount of a compound as defined in any of one Claims 1 to 32.
46. A method of stabilising a DNA duplex formed between first and second single strands of DNA, which method comprises contacting that DNA duplex with a compound as defined in any of one Claims 1 to 32.
47. A method of enhancing the difference in melting temperatures between first and second DNA duplexes, wherein each DNA duplex is formed from a first single strand of DNA that is the same in each duplex and a second single strand of DNA that is different in each duplex, which

method comprises contacting each DNA duplex with a compound as defined in any of one Claims 1 to 32.


48. A process for the preparation of compounds of formula I as defined in Claim 16 which comprises:

(a) reaction of a compound of formula III,




wherein A^a represents A or, when a represents 0, then A^a may also represent A^2 and Q, D, A and A^2 are as defined in Claim 16 and a is as defined below,

with a compound of formula IV,




wherein A^b represents a direct bond or $-\text{A}^1-\text{C}(\text{O})-$, as appropriate, L^1 represents a leaving group, a and b both represent integers from 0 to 5, the sum of the two being 2, 3, 4 or 5, and R^1 and Q are as defined in Claim 16;

(b) for compounds of formula I in which R^1 represents $\text{D}-\text{A}-\text{N}(\text{H})-[\text{Q}]_n-\text{C}(\text{O})-\text{E}-\text{C}(\text{O})-$, reaction of two equivalents of a compound of formula V,



wherein Q, n, A and D are as defined in Claim 16, with a compound of formula VI,



wherein L^2 represents a leaving group, the two L^2 groups being the same or different, and E is as defined in Claim 16; or

(c) deprotection of a protected derivative of a compound of formula I as defined in Claim 16.

49. A compound of formula V, as defined in Claim 48, or a protected derivative thereof.

1/9

CE Conditions

Capillary Electrophoresis: TSP CE1000

Capillary (type/length): Bare fused silica (375 μ m O.D., 50 μ m I.D.) effective 32cm. total 40cm.

Applied Voltage: 25kV

Injection: Hydrodynamic, 5sec.

Capillary temperature 20°C \pm 0.1°C

Buffer: 0.02M Borate buffer, pH7.5



UV detection: 260nm.

DNA Cone: 20 μ M

Key: Peak 1: AAATTATATTAT (AT)

Peak 2: GGGCCGCGCCGC (GC)

Peak 3: Netropsin:AT complex

**Netropsin**

Competitive binding studies with netropsin.

Fig 1

2/9

CE Conditions

Capillary Electrophoresis: Agilent G1600

Capillary (type/length): Bare fused silica (375 μ m O.D., 50 μ m I.D.) effective 25cm. total 33.5cm.

Applied Voltage: 25kV

Injection: Pressure, (50mBar, 5sec)


Capillary temperature 20°C \pm 0.1°C

Buffer: 0.02M Borate buffer, pH7.5


UV detection: 260nm.

DNA Cone: 20 μ M

Key: Peak 1: AAATTATATTAT (AT)
 Peak 2: GGGCCGCGCCGC (GC)
 Peak 3: 1241:AT complex



12/41



Competitive binding studies with compound 12/41

Fig.2

3/9

CE Conditions

Capillary Electrophoresis: Beckman P/ACE MDQ.

Capillary (type/length): Bare fused SILICA(375 μ mO.D., 50 μ m I.D., effective 21cm, total 31.2 cm.

Applied Voltage: 25kV

Injection: Pressure, (20 P.S.I., 0.5 sec)

Capillary temperature 20°C \pm 0.1°C


Buffer: 0.02M Borate buffer, pH7.5

UV detection: 260nm.


DNA Cone: 20 μ M

Key: Peak 1: CGACTAGTCG (ACTA)

Peak 2: 1320:ACTA complex



13/20



Capillary electrophoresis studies of the interaction between compound 13/20 and DNA decamer CGACTAGTCG.

Fig.3

4/9

CE Conditions

Capillary Electrophoresis: Beckman P/ACE MDQ.

Capillary (type/length): Bare fused SILICA(375 μ m O.D., 50 μ m I.D.yeffective 21 cm, total 31.2 cm.

Applied Voltage: 25kV

Injection: Pressure, (20 P.S.I., 0.5 sec)



Capillary temperature 20°C \pm 0.1°C

Buffer: 0.02M Borate buffer, pH7.5

UV detection: 260nm.

DNA Cone: 20 μ M

Key: Peak 1: CGACTAGTCG (ACTA)

**Distamycin A**

Capillary electrophoresis studies of the interaction between
distamycin and DNA decamer CGACTAGTCG.

Fig.4

5/9

CE Conditions

Capillary Electrophoresis: Agilent G1600

Capillary (type/length): Bare fused SILICA(375 μ m)

O.D., 50 μ m I.D.yeffective 25cm, total 33.5 cm.

Applied Voltage: 25kV

Injection: Pressure, (50mBar, 5 sec)

Capillary temperature $20^{\circ}\text{C} \pm 0.1^{\circ}\text{C}$

Buffer: 0.02M Borate buffer, pH7.5

UV detection: 260nm.

DNA Cone: 20 μ M


Key: Peak 1: AAATTATATTAT (AT).

Peak 2: CGACTAGTCG (ACTA).


Peak 3: 1320: ACTA complex.

Peak 4: 1320: AT complex.

* Ratio with respect to each duplex present



13/20



Competitive binding studies with compound 13/20

Fig 5

6/9

CE Conditions

Capillary Electrophoresis: Beckman P/ACE MDQ.

Capillary (type/length): Bare fused Silica(375 μ m O.D., 50 μ m I.D.) effective 21cm, total 31.2 cm.

Applied Voltage: 25kV

Injection: Pressure, (20 P.S.I., 0.5 sec)


Capillary temperature 20°C \pm 0.1°C

Buffer: 0.02M Borate buffer, pH7.5


UV detection: 260nm.

DNA Cone: 20 μ M

Key: Peak 1: CGACTAGTGG (ACTA2)
Peak 1: 1320: ACTA2 complex



1320



Capillary electrophoresis studies of the interaction between compound 13/20 and DNA decamer CCACTAGTGG

Fig. 6

7/9

Footprinting studies with compound 13/20




Fig. 7

5'-GGATCCATATGCGGCAATACACATGGCCGATTTCCTCAACTGCACTAGTCGTAGCGGATCAAGGTTAAGCTCCCGTTCTATCC
 x XXXXXXxxxx

TGGTATAGCAATTAGGGCGTGAAAGAGTTATGTAAAGTACGTCCGGTGGGGTCTGTTTGTGTCATCTCAGCCTCGAATGCGGATC

(Experiment carried out at 0.03 μ M of compound 13/20)

Footprinting studies with compound 13/51




Fig. 8

5'-GGATCCATATGCGGCAATACACATGGCCGATTTCCAAGTCAAGTTAAGTCCCGTTCTATCC
xxxxx(0.1)

xxxx(3)	xxxxxxx(0.1)	xxxxx(1)	xxxxxx(0.01)	xxxxx(0.1)
TGGTATAGCAATTAGGGCGTGAAGAGTTATGTAAAGTACGTCCGGTGGGCTCTGTTTTGTCACTCAGCCCTCGAATGCGGATC				

The numbers in brackets refer to the concentration of compound 13/51 (in μM) at which the relevant footprint was observed.

Effect of DNA Minor Groove Binding Compounds upon LPS-Stimulated NF κ B-DNA Binding Activity in Murine Macrophages




Fig. 9

C= control; n.s. = non-specific